

## FP3 Numerical Methods for the Solution of First Order Differential Equations Questions

- 5 (a) The function  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = x \ln x + \frac{y}{x}$

and  $y(1) = 1$

- (i) Use the Euler formula

$$y_{r+1} = y_r + h f(x_r, y_r)$$

with  $h = 0.1$ , to obtain an approximation to  $y(1.1)$ . *(3 marks)*

- (ii) Use the formula

$$y_{r+1} = y_{r-1} + 2h f(x_r, y_r)$$

with your answer to part (a)(i) to obtain an approximation to  $y(1.2)$ , giving your answer to three decimal places. *(4 marks)*

- (b) (i) Show that  $\frac{1}{x}$  is an integrating factor for the first-order differential equation

$$\frac{dy}{dx} - \frac{1}{x}y = x \ln x \quad (3 \text{ marks})$$

- (ii) Solve this differential equation, given that  $y = 1$  when  $x = 1$ . *(6 marks)*
- (iii) Calculate the value of  $y$  when  $x = 1.2$ , giving your answer to three decimal places. *(1 mark)*
-

2 The function  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = \frac{x^2 + y^2}{xy}$

and  $y(1) = 2$

(a) Use the Euler formula

$$y_{r+1} = y_r + hf(x_r, y_r)$$

with  $h = 0.1$ , to obtain an approximation to  $y(1.1)$ . (3 marks)

(b) Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where  $k_1 = hf(x_r, y_r)$  and  $k_2 = hf(x_r + h, y_r + k_1)$  and  $h = 0.1$ , to obtain an approximation to  $y(1.1)$ , giving your answer to four decimal places. (6 marks)

---

1 The function  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = \ln(1 + x^2 + y)$

and  $y(1) = 0.6$

(a) Use the Euler formula

$$y_{r+1} = y_r + hf(x_r, y_r)$$

with  $h = 0.05$ , to obtain an approximation to  $y(1.05)$ , giving your answer to four decimal places. (3 marks)

(b) Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where  $k_1 = hf(x_r, y_r)$  and  $k_2 = hf(x_r + h, y_r + k_1)$  and  $h = 0.05$ , to obtain an approximation to  $y(1.05)$ , giving your answer to four decimal places. (6 marks)

---

2 The function  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = \sqrt{x^2 + y^2 + 3}$

and  $y(1) = 2$

(a) Use the Euler formula

$$y_{r+1} = y_r + hf(x_r, y_r)$$

with  $h = 0.1$ , to obtain an approximation to  $y(1.1)$ , giving your answer to four decimal places. *(3 marks)*

(b) Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where  $k_1 = hf(x_r, y_r)$  and  $k_2 = hf(x_r + h, y_r + k_1)$  and  $h = 0.1$ , to obtain an approximation to  $y(1.1)$ , giving your answer to four decimal places. *(6 marks)*

---