

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS**

**Advanced Subsidiary General Certificate of Education  
Advanced General Certificate of Education**

**MATHEMATICS**

**4726**

**Further Pure Mathematics 2**

Tuesday                      **6 JUNE 2006**                      Afternoon                      1 hour 30 minutes

Additional materials:  
8 page answer booklet  
Graph paper  
List of Formulae (MF1)

**TIME**    1 hour 30 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

---

**This question paper consists of 4 printed pages.**

- 1 Find the first three non-zero terms of the Maclaurin series for

$$(1 + x) \sin x,$$

simplifying the coefficients.

[3]

- 2 (i) Given that  $y = \tan^{-1} x$ , prove that  $\frac{dy}{dx} = \frac{1}{1+x^2}$ .

[3]

- (ii) Verify that  $y = \tan^{-1} x$  satisfies the equation

$$(1+x^2) \frac{d^2y}{dx^2} + 2x \frac{dy}{dx} = 0.$$

[3]

- 3 The equation of a curve is  $y = \frac{x+1}{x^2+3}$ .

- (i) State the equation of the asymptote of the curve.

[1]

- (ii) Show that  $-\frac{1}{6} \leq y \leq \frac{1}{2}$ .

[5]

- 4 (i) Using the definition of  $\cosh x$  in terms of  $e^x$  and  $e^{-x}$ , prove that

$$\cosh 2x = 2 \cosh^2 x - 1.$$

[3]

- (ii) Hence solve the equation

$$\cosh 2x - 7 \cosh x = 3,$$

giving your answer in logarithmic form.

[4]

- 5 (i) Express  $t^2 + t + 1$  in the form  $(t+a)^2 + b$ .


[1]

- (ii) By using the substitution  $\tan \frac{1}{2}x = t$ , show that

$$\int_0^{\frac{1}{2}\pi} \frac{1}{2 + \sin x} dx = \frac{\sqrt{3}}{9} \pi.$$

[6]

6



The diagram shows the curve with equation  $y = 3^x$  for  $0 \leq x \leq 1$ . The area  $A$  under the curve between these limits is divided into  $n$  strips, each of width  $h$  where  $nh = 1$ .

(i) By using the set of rectangles indicated on the diagram, show that  $A > \frac{2h}{3^h - 1}$ . [3]

(ii) By considering another set of rectangles, show that  $A < \frac{(2h)3^h}{3^h - 1}$ . [3]

(iii) Given that  $h = 0.001$ , use these inequalities to find values between which  $A$  lies. [2]

7 The equation of a curve, in polar coordinates, is

$$r = \sqrt{3} + \tan \theta, \quad \text{for } -\frac{1}{3}\pi \leq \theta \leq \frac{1}{4}\pi.$$

(i) Find the equation of the tangent at the pole. [2]

(ii) State the greatest value of  $r$  and the corresponding value of  $\theta$ . [2]

(iii) Sketch the curve. [2]

(iv) Find the exact area of the region enclosed by the curve and the lines  $\theta = 0$  and  $\theta = \frac{1}{4}\pi$ . [5]

8 The curve with equation  $y = \frac{\sinh x}{x^2}$ , for  $x > 0$ , has one turning point.

(i) Show that the  $x$ -coordinate of the turning point satisfies the equation  $x - 2 \tanh x = 0$ . [3]

(ii) Use the Newton-Raphson method, with a first approximation  $x_1 = 2$ , to find the next two approximations,  $x_2$  and  $x_3$ , to the positive root of  $x - 2 \tanh x = 0$ . [5]

(iii) By considering the approximate errors in  $x_1$  and  $x_2$ , estimate the error in  $x_3$ . (You are not expected to evaluate  $x_4$ .) [3]

[Question 9 is printed overleaf.]

9 (i) Given that  $y = \sinh^{-1} x$ , prove that  $y = \ln(x + \sqrt{x^2 + 1})$ . [3]

(ii) It is given that, for non-negative integers  $n$ ,

$$I_n = \int_0^\alpha \sinh^n \theta \, d\theta,$$

where  $\alpha = \sinh^{-1} 1$ . Show that

$$nI_n = \sqrt{2} - (n-1)I_{n-2}, \quad \text{for } n \geq 2. \quad [6]$$

(iii) Evaluate  $I_4$ , giving your answer in terms of  $\sqrt{2}$  and logarithms. [4]