

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS**

**Advanced Subsidiary General Certificate of Education  
Advanced General Certificate of Education**

**MATHEMATICS**

**4730**

**Mechanics 3**

Monday                      **22 MAY 2006**                      Morning                      1 hour 30 minutes

Additional materials:  
8 page answer booklet  
Graph paper  
List of Formulae (MF1)

**TIME**    1 hour 30 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- The acceleration due to gravity is denoted by  $g \text{ m s}^{-2}$ . Unless otherwise instructed, when a numerical value is needed, use  $g = 9.8$ .
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**


---

**This question paper consists of 4 printed pages.**

- 1 A ball of mass  $0.4 \text{ kg}$  is moving in a straight line, with speed  $25 \text{ m s}^{-1}$ , when it is struck by a bat. The bat exerts an impulse of magnitude  $20 \text{ N s}$  and the ball is deflected through an angle of  $90^\circ$ . Calculate
- (i) the direction of the impulse, [3]
- (ii) the speed of the ball immediately after it is struck. [3]

- 2 A duck of mass  $2 \text{ kg}$  is travelling with horizontal speed  $4 \text{ m s}^{-1}$  when it lands on a lake. The duck is brought to rest by the action of resistive forces, acting in the direction opposite to the duck's motion and having total magnitude  $(2v + 3v^2) \text{ N}$ , where  $v \text{ m s}^{-1}$  is the speed of the duck. Show that the duck comes to rest after travelling approximately  $1.30 \text{ m}$  from the point of its initial contact with the surface of the lake. [8]

3




Two uniform rods  $AB$  and  $AC$ , of equal lengths, and of weights  $200 \text{ N}$  and  $360 \text{ N}$  respectively, are freely jointed at  $A$ . The mid-points of the rods are joined by a taut light inextensible string. The rods are in equilibrium in a vertical plane with  $B$  and  $C$  in contact with a smooth horizontal surface. The point  $A$  is  $2.1 \text{ m}$  above the surface and  $BC = 1.4 \text{ m}$  (see diagram).

- (i) Show that the force exerted on  $AB$  at  $B$  has magnitude  $240 \text{ N}$  and find the tension in the string. [6]
- (ii) Find the horizontal and vertical components of the force exerted on  $AB$  at  $A$  and state their directions. [3]
- 4 A particle is connected to a fixed point by a light inextensible string of length  $2.45 \text{ m}$  to make a simple pendulum. The particle is released from rest with the string taut and inclined at  $0.1$  radians to the downward vertical.
- (i) Show that the motion of the particle is approximately simple harmonic with period  $3.14 \text{ s}$ , correct to 3 significant figures. [5]

Calculate, in either order,

- (ii) the angular speed of the pendulum when it has moved  $0.04$  radians from the **initial** position, [3]
- (iii) the time taken by the pendulum to move  $0.04$  radians from the **initial** position. [3]

5



Two uniform smooth spheres  $A$  and  $B$ , of equal radius, have masses 2 kg and 3 kg respectively. They are moving on a horizontal surface when they collide. Immediately before the collision  $A$  is moving with speed  $12 \text{ m s}^{-1}$  at  $60^\circ$  to the line of centres, and  $B$  is moving with speed  $8 \text{ m s}^{-1}$  along the line of centres (see diagram). The coefficient of restitution between the spheres is 0.5. Find the speed and direction of motion of each sphere after the collision. [12]

- 6 A bungee jumper of mass 70 kg is joined to a fixed point  $O$  by a light elastic rope of natural length 30 m and modulus of elasticity 1470 N. The jumper starts from rest at  $O$  and falls vertically. The jumper is modelled as a particle and air resistance is ignored.
- (i) Find the distance fallen by the jumper when maximum speed is reached. [4]
- (ii) Show that this maximum speed is  $26.9 \text{ m s}^{-1}$ , correct to 3 significant figures. [4]
- (iii) Find the extension of the rope when the jumper is at the lowest position. [4]

[Question 7 is printed overleaf.]




Fig. 1




Fig. 2

A smooth horizontal cylinder of radius 0.6 m is fixed with its axis horizontal and passing through a fixed point  $O$ . A light inextensible string of length  $0.6\pi$  m has particles  $P$  and  $Q$ , of masses 0.3 kg and 0.4 kg respectively, attached at its ends. The string passes over the cylinder and is held at rest with  $P$ ,  $O$  and  $Q$  in a straight horizontal line (see Fig. 1). The string is released and  $Q$  begins to descend. When the line  $OP$  makes an angle  $\theta$  radians,  $0 \leq \theta \leq \frac{1}{2}\pi$ , with the horizontal, the particles have speed  $v$  m s<sup>-1</sup> (see Fig. 2).

- (i) By considering the total energy of the system, or otherwise, show that

$$v^2 = 6.72\theta - 5.04 \sin \theta. \quad [5]$$

- (ii) Show that the magnitude of the contact force between  $P$  and the cylinder is

$$(5.46 \sin \theta - 3.36\theta) \text{ newtons.}$$

Hence find the value of  $\theta$  for which the magnitude of the contact force is greatest. [6]

- (iii) Find the transverse component of the acceleration of  $P$  in terms of  $\theta$ . [3]