

# GCE 2005

## *January Series*



ASSESSMENT and  
QUALIFICATIONS  
ALLIANCE

# Mark Scheme

## Mathematics

MM1B

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website:  
[www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2005 AQA and its licensors. All rights reserved.

#### COPYRIGHT


AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

*Dr Michael Cresswell Director G*

## MM1B

Q	Solution	Marks	Total	Comments		
1(a)(i)	$40 = 12 + 100a$	M1	2	Use of a constant acceleration equation to form equation for $a$ AG; correct answer from correct working		
	$a = \frac{40-12}{100} = 0.28 \text{ ms}^{-2}$ AG	A1				
	(ii)	$s = \frac{1}{2}(12 + 40) \times 100$	M1	2	Expression for distance, using $t = 100$ Correct final distance	
$= 2600 \text{ m}$		A1				
(c)	$F - 40000 = 200 \times 1000 \times 0.28$	M1	3	Three term equation of motion Correct equation Correct force		
	$F = 40000 + 56000 = 96000 \text{ N}$	A1				
		A1				
<b>Total</b>			<b>7</b>			
2(a)	$12 \begin{bmatrix} 4 \\ 7 \end{bmatrix} + 4 \begin{bmatrix} 2 \\ 3 \end{bmatrix} = 16\mathbf{v}$	M1 A1	4	Three term momentum equation Correct equation  Solving for $\mathbf{v}$ Correct velocity		
	$\mathbf{v} = \frac{1}{16} \begin{bmatrix} 56 \\ 96 \end{bmatrix} = \begin{bmatrix} 3.5 \\ 6.0 \end{bmatrix} \text{ ms}^{-1}$	m1 A1				
(b)	$12 \begin{bmatrix} 4 \\ 7 \end{bmatrix} + 4\mathbf{u} = 16 \begin{bmatrix} 1 \\ 4 \end{bmatrix}$	M1 A1			3	Three term momentum equation Correct equation  Correct velocity
	$\mathbf{u} = \frac{1}{4} \begin{bmatrix} -32 \\ -20 \end{bmatrix} = \begin{bmatrix} -8 \\ -5 \end{bmatrix} \text{ ms}^{-1}$	A1				
<b>Total</b>			<b>7</b>			
3 (a)		B1	1	Correct diagram		
(b)	$40 \cos 30^\circ - F = 25 \times 0.1$	M1	3	Three term equation of motion Correct equation AG; correct force from correct working		
	$F = 40 \cos 30^\circ - 2.5 = 32.1 \text{ N}$	A1 A1				
(c)	$R + 40 \sin 30^\circ = 25 \times 9.8$	M1	3	Resolving vertically Correct equation AG; correct force from correct working		
	$R = 225 \text{ N}$	A1 A1				
(d)	$32.1 = 225\mu$	M1	2	use of $F = \mu R$  Correct $\mu$		
	$\mu = \frac{32.1}{225} = 0.143$	A1				
(e)	Friction will decrease as normal reaction decreases	B1 B1	2	Decrease in friction Normal reaction decreases		
<b>Total</b>			<b>11</b>			

**MM1B (cont)**

Q	Solution	Marks	Total	Comments
4(a)	Light or smooth	B1	1	Acceptable assumption
(b)	$5g - T = 5a$ $T - 2g = 2a$ $3g = 7a$ $a = \frac{3g}{7} = 4.2 \text{ ms}^{-2}$	M1 A1 M1 A1 A1	5	Three term equation of motion for one particle Correct equation Three term equation of motion for other particle Correct equation AG; correct acceleration from correct working
(c)	$T = 2 \times 4.2 + 2 \times 9.8 = 28 \text{ N}$	M1 A1	2	Substitute $a = 4.2$ into one equation of motion Correct tension
<b>Total</b>			<b>8</b>	
5(a)	$200 \sin 30^\circ = T \sin 45^\circ$ $T = \frac{200 \sin 30^\circ}{\sin 45^\circ} = 141 \text{ N}$	M1 A1 A1	3	Resolving horizontally Correct equation AG; correct $T$ from correct working
(b)	$200 \cos 30^\circ + 141 \cos 45^\circ + R = 500 \times 9.8$ $R = 4630 \text{ N}$	M1 A1 A1 A1	4	Resolving vertically with four terms Correct values Correct signs Correct $R$
<b>Total</b>			<b>7</b>	
6(a)	$\frac{\sin 60^\circ}{6} = \frac{\sin \alpha}{2}$ $\alpha = 16.8^\circ$	M1 A1 A1 A1	4	Use of sine rule Correct LHS Correct RHS AG; correct $\alpha$ from correct working
(b)	$\frac{v}{\sin(180 - 60 - 16.8)} = \frac{6}{\sin 60^\circ}$ $v = 6.74 \text{ or } 6.75 \text{ ms}^{-1}$	M1 A1 A1	3	use of sine rule to find $v$ Correct equation Correct $v$
<b>Total</b>			<b>7</b>	

MM1B (cont)

Q	Solution	Marks	Total	Comments
7 (a)	$-\mathbf{i} + \mathbf{j} = 2\mathbf{i} - \mathbf{j} + 10\mathbf{a}$ $\mathbf{a} = -0.3\mathbf{i} + 0.2\mathbf{j}$	M1 A1 A1	3	Use of velocity equation Correct equation Correct $\mathbf{a}$
(b)	$\mathbf{r} = (2\mathbf{i} - \mathbf{j})t + \frac{1}{2}(-0.3\mathbf{i} + 0.2\mathbf{j})t^2 + 20\mathbf{i}$ $= (2t - 0.15t^2 + 20)\mathbf{i} + (-t + 0.1t^2)\mathbf{j}$	M1 A1 A1 ft	3	Use of constant acceleration equation for position Correct $\mathbf{i}$ component Correct $\mathbf{j}$ component ft incorrect acceleration
(c) (i)	$\mathbf{r}(20) = (2 \times 20 - 0.15 \times 20^2 + 20)\mathbf{i} + (-20 + 0.1 \times 20^2)\mathbf{j}$ $= 0\mathbf{i} + 20\mathbf{j}$ so due north of origin	M1 A1	2	Substituting $t = 20$ into their expression for $\mathbf{r}$ Correct conclusion from correct working
(c)(ii)	$\mathbf{v}(20) = 2\mathbf{i} - \mathbf{j} + 20(-0.3\mathbf{i} + 0.2\mathbf{j})$ $= -4\mathbf{i} + 3\mathbf{j}$ $v(20) = \sqrt{4^2 + 3^2} = 5 \text{ ms}^{-1}$	M1 A1 m1 A1ft	4	Finding velocity at $t = 20$ Correct velocity Finding magnitude Correct speed ft incorrect acceleration
<b>Total</b>			<b>12</b>	
8(a)	Ball is a particle No air resistance	B1 B1	2	One appropriate assumption Second appropriate assumption
(b)(i)	$0 = 12 \sin 40^\circ - 9.8t$ $t = \frac{12 \sin 40^\circ}{9.8} = 0.787 \text{ s}$	M1 A1 M1 A1	4	Equation to find time at maximum height Correct equation Solving for $t$ Correct time
(ii)	$h = 12 \sin 40^\circ \times 0.7871 - 4.9 \times 0.7871^2$ $= 3.04 \text{ m}$	M1 A1 A1	3	Substituting time from previous into expression for height Correct expression AG; correct height from correct working
(c)	$2.44 = 12 \sin 40^\circ t - 4.9t^2$ $4.9t^2 - 12 \sin 40^\circ t + 2.44 = 0$ $t = 0.4385 \text{ or } 1.136$ $s = 12 \cos 40^\circ \times 1.136 = 10.4 \text{ m}$	M1 A1 A1 m1 A1 M1 A1	7	Equation for time to get to the bar, based on height being 2.44 Correct LHS Correct RHS Solving quadratic Correct time / times Substituting their larger time into an expression for the horizontal displacement Correct distance
<b>Total</b>			<b>16</b>	
<b>TOTAL</b>			<b>75</b>	