



General Certificate of Education

Mathematics 6360

MFP1 Further Pure 1

Mark Scheme

2005 examination – June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous		
	incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	OE	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

Application of Mark Scheme

No method shown:

Correct answer without working
Incorrect answer without working

mark as in scheme
zero marks unless specified otherwise

More than one method / choice of solution:

2 or more complete attempts, neither/none crossed out

mark both/all fully and award the mean
mark rounded down

1 complete and 1 partial attempt, neither crossed out

award credit for the complete solution only

Crossed out work

do not mark unless it has not been replaced

Alternative solution using a correct or partially correct method

award method and accuracy marks as
appropriate

MFP1

Q	Solution	Marks	Total	Comments
1(a)(i)	$\mathbf{A} + \mathbf{B} = \begin{bmatrix} 3 & 6 \\ 6 & 3 \end{bmatrix}$	M1A1	2	M1A0 if 3 entries correct
(ii)	$\mathbf{AB} = \begin{bmatrix} 8 & 6 \\ 6 & 8 \end{bmatrix}$	M1A1	2	Ditto
(b)	$\mathbf{A} + \mathbf{B} - \mathbf{AB} = \begin{bmatrix} -5 & 0 \\ 0 & -5 \end{bmatrix}$	B1F		ft wrong answers in (a)
	$\dots = -5 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$	B1	2	
	Total		6	
2	$x = 0.5 \Rightarrow y' = \sin 1$ First increment is $0.1 \sin 1$ $x = 0.6 \Rightarrow y \approx 1.084$ Second increment is $0.1 \sin 1.2$ $x = 0.7 \Rightarrow y \approx 1.177 \approx 1.18$	M1 m1A1 A1 m1 A1F	6	Max 4/6 if degrees used PI by correct answer at end Ft error in $y(0.6)$
	Total		6	
3(a)	$\Sigma r^2(r-1) = \Sigma r^3 - \Sigma r^2$ Good progress with expansion Factors n and $n + 1$ found $\dots = \frac{1}{12}n(n^2 - 1)(3n + 2)$	M1 m1 A1 A1	4	With attempt to use the given formulae or use of common factors Allow verification here Convincingly shown (AG)
(b)	Use of $f(11) - f(3)$ in above $f(11) = 3850$ $f(3) = 22$ (so answer is 3828)	M1 A1 A1	3	M1 for $f(11) - f(4)$ PI by correct answer ditto
	Total		7	
4(a)	$(2 + h)^3 = 8 + ah + bh^2 + h^3$ $(2 + h)^3 = 8 + 12h + 6h^2 + h^3$ $f(2 + h) - f(2) = 13h + 6h^2 + h^3$	M1 A1A1 m1A1F	5	A1 for each of a, b ; PI Ft one coeff. wrong
(b)	Divide by h and let $h \rightarrow 0$ $f'(2) = p = 13$	M1 A1F	2	NMS B1F ft wrong value of p
	Total		7	

MFP1 (cont)

Q	Solution	Mark	Total	Comments
5(a)	Use of $\tan \frac{\pi}{3} = \sqrt{3}$ GS is $\frac{1}{3}(\frac{\pi}{3} + n\pi)$	M1 m1A1	3	Degrees used - 1-mark penalty m1A0 for $\frac{\pi}{9} + n\pi$ or for correct answer plus extra solutions
(b)	One solution is 0 GS is $\frac{1}{3}n\pi$	M1A1 m1A1F	4	OE m1A0 for $n\pi$ or for correct answer plus extra solns; ft wrong first solution
Total			7	
6(a)(i)	$\alpha + \beta = 4, \alpha\beta = 13$	B1B1	2	convincingly shown (AG)
(ii)	$\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$... = $4^2 - 26 = -10$	M1 A1	2	
(iii)	The square of a real number is positive (or zero) The sum of two such squares is positive (or zero)	E1 E1	2	
(b)(i)	$(\alpha + i) + (\beta + i) = 4 + 2i$	B1F	1	
(ii)	$(\alpha + i)(\beta + i) = 12 + 4i$	M1A1F	2	ditto
(c)	Correct coeff of x or constant term $x^2 - (4 + 2i)x + (12 + 4i) = 0$	M1 A1F	2	Using c's answers in (b) ft wrong answers in (b)
Total			11	
7(a)(i)	$D(4, 0)$ $E(8, -4), F(10, -2)$	M1A1 A1A1	4	M1 if at least one point correct Ft one error NMS 2/2; 1/2 for AWRT 2.8 NMS 2/2; condone 315° ; 1/2 for AWRT $44-46^\circ$ OE
(ii)	Correct sketch	m1A1F	2	
(b)(i)	Scale factor is $2\sqrt{2}$	M1A1	2	
(ii)	Angle 45°	M1A1	2	
Total			10	
8(a)	P is $(2, 0)$	B1	1	ft wrong value for x_P ft numerical error
(b)	PQ is $y = 2(x - 2)$ Elimination of y (or of x) $(x - 2)(5x - 22) = 0$ Q is $(4.4, 4.8)$	M1A1F m1A1F A1 A1A1	7	
Total			8	
9(a)(i)	Asymptote is $y = 1$	B1	1	
(ii)	Denominator never zero	E2,1	2	
(b)	$f(x) = k \Rightarrow (1 - k)x^2 + 4x - 9k = 0$ Equal roots if $16 + 36k(1 - k) = 0$ ie if $9k^2 - 9k - 4 = 0$	M1A1 m1 A1	4	
(c)	Solving quadratic for k $k = -\frac{1}{3}$ or $k = \frac{4}{3}$	M1 A1		
	$4x^2 + 12x + 9 = 0$ or $x^2 - 12x + 36 = 0$	m1		
	SPs when $x = -\frac{3}{2} \dots$; and when $x = 6 \dots$	A1 A1		
	SPs are $(-\frac{3}{2}, -\frac{1}{3})$ and $(6, \frac{4}{3})$	A1	6	
Total			13	
Total			75	