

MATHEMATICS
Unit Further Pure 1

MFP1

Thursday 15 January 2009 9.00 am to 10.30 am

For this paper you must have:

- an 8-page answer book
- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MFP1.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

Information

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

Answer **all** questions.

- 1 A curve passes through the point $(0, 1)$ and satisfies the differential equation

$$\frac{dy}{dx} = \sqrt{1 + x^2}$$

Starting at the point $(0, 1)$, use a step-by-step method with a step length of 0.2 to estimate the value of y at $x = 0.4$. Give your answer to five decimal places. *(5 marks)*

- 2 The complex number $2 + 3i$ is a root of the quadratic equation

$$x^2 + bx + c = 0$$

where b and c are real numbers.

- (a) Write down the other root of this equation. *(1 mark)*
- (b) Find the values of b and c . *(4 marks)*

- 3 Find the general solution of the equation

$$\tan\left(\frac{\pi}{2} - 3x\right) = \sqrt{3} \quad (5 \text{ marks})$$

- 4 It is given that

$$S_n = \sum_{r=1}^n (3r^2 - 3r + 1)$$

- (a) Use the formulae for $\sum_{r=1}^n r^2$ and $\sum_{r=1}^n r$ to show that $S_n = n^3$. *(5 marks)*

- (b) Hence show that $\sum_{r=n+1}^{2n} (3r^2 - 3r + 1) = kn^3$ for some integer k . *(2 marks)*

5 The matrices \mathbf{A} and \mathbf{B} are defined by

$$\mathbf{A} = \begin{bmatrix} k & k \\ k & -k \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} -k & k \\ k & k \end{bmatrix}$$

where k is a constant.

(a) Find, in terms of k :

(i) $\mathbf{A} + \mathbf{B}$; *(1 mark)*

(ii) \mathbf{A}^2 . *(2 marks)*

(b) Show that $(\mathbf{A} + \mathbf{B})^2 = \mathbf{A}^2 + \mathbf{B}^2$. *(4 marks)*

(c) It is now given that $k = 1$.

(i) Describe the geometrical transformation represented by the matrix \mathbf{A}^2 . *(2 marks)*

(ii) The matrix \mathbf{A} represents a combination of an enlargement and a reflection. Find the scale factor of the enlargement and the equation of the mirror line of the reflection. *(3 marks)*

6 A curve has equation

$$y = \frac{(x-1)(x-3)}{x(x-2)}$$

(a) (i) Write down the equations of the three asymptotes of this curve. *(3 marks)*

(ii) State the coordinates of the points at which the curve intersects the x -axis. *(1 mark)*

(iii) Sketch the curve.


(You are given that the curve has no stationary points.) *(4 marks)*

(b) Hence, or otherwise, solve the inequality

$$\frac{(x-1)(x-3)}{x(x-2)} < 0 \quad \text{span style="float: right;">*(2 marks)*$$

Turn over ►

- 7 The points $P(a, c)$ and $Q(b, d)$ lie on the curve with equation $y = f(x)$. The straight line PQ intersects the x -axis at the point $R(r, 0)$. The curve $y = f(x)$ intersects the x -axis at the point $S(\beta, 0)$.



- (a) Show that

$$r = a + c \left(\frac{b - a}{c - d} \right) \quad (4 \text{ marks})$$

- (b) Given that

$$a = 2, b = 3 \text{ and } f(x) = 20x - x^4$$

- (i) find the value of r ; (3 marks)
- (ii) show that $\beta - r \approx 0.18$. (3 marks)

- 8 For each of the following improper integrals, find the value of the integral **or** explain why it does not have a value:

(a) $\int_1^{\infty} x^{-\frac{3}{4}} dx;$ (3 marks)

(b) $\int_1^{\infty} x^{-\frac{5}{4}} dx;$ (3 marks)

(c) $\int_1^{\infty} (x^{-\frac{3}{4}} - x^{-\frac{5}{4}}) dx.$ (1 mark)

- 9 A hyperbola H has equation

$$x^2 - \frac{y^2}{2} = 1$$

- (a) Find the equations of the two asymptotes of H , giving each answer in the form $y = mx$. (2 marks)

- (b) Draw a sketch of the two asymptotes of H , using roughly equal scales on the two coordinate axes. Using the same axes, sketch the hyperbola H . (3 marks)

- (c) (i) Show that, if the line $y = x + c$ intersects H , the x -coordinates of the points of intersection must satisfy the equation

$$x^2 - 2cx - (c^2 + 2) = 0$$
 (4 marks)

- (ii) Hence show that the line $y = x + c$ intersects H in two distinct points, whatever the value of c . (2 marks)

- (iii) Find, in terms of c , the y -coordinates of these two points. (3 marks)

END OF QUESTIONS

There are no questions printed on this page

There are no questions printed on this page

There are no questions printed on this page