



General Certificate of Education

Mathematics 6360

MFP2 Further Pure 2

Mark Scheme

2006 examination – January series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Key To Mark Scheme And Abbreviations Used In Marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MFP2

Q	Solution	Marks	Total	Comments
1(a)	$\frac{1}{r^2} - \frac{1}{(r+1)^2} = \frac{(r+1)^2 - r^2}{r^2(r+1)^2}$ $= \frac{2r+1}{r^2(r+1)^2}$	M1 A1	2	AG
(b)	$\frac{3}{1^2 \times 2^2} = \frac{1}{1^2} - \frac{1}{2^2}$ $\frac{5}{2^2 \times 3^2} = \frac{1}{2^2} - \frac{1}{3^2}$ $\frac{7}{3^2 \times 4^2} = \frac{1}{3^2} - \frac{1}{4^2}$ $\frac{2n+1}{n^2(n+1)^2} = \frac{1}{n^2} - \frac{1}{(n+1)^2}$ <p>Clear cancellation</p> $1 - \frac{1}{(n+1)^2}$	M1A1 M1	4	A1 for at least 3 lines
Total			6	
2(a)	$p = -4$ $(\alpha + \beta + \gamma)^2 = \sum \alpha^2 + 2 \sum \alpha\beta$ $16 = 20 + 2 \sum \alpha\beta$ $\sum \alpha\beta = -2$ $q = -2$	B1 M1 A1 A1F A1F	5	
(b)	$3 - i$ is a root Third root is -2 $\alpha\beta\gamma = (3+i)(3-i)(-2)$ $= -20$ $r = +20$	B1 B1F M1 A1F A1F	5	Real $\alpha\beta\gamma$ Real r
Alternative to (b)				
	Substitute $3 + i$ into equation $(3 + i)^2 = 8 + 6i$ $(3 + i)^3 = 18 + 26i$ $r = 20$	M1 B1 B1 A2,1,0		Provided r is real
Total			10	


MFP2 (cont)

Q	Solution	Marks	Total	Comments
3(a)	$\frac{1+i}{1-i} = \frac{(1+i)^2}{1-i^2} = i$	M1A1	2	AG
(b)	$ z_2 = \sqrt{\frac{1}{4} + \frac{3}{4}} = 1 = z_1 $	M1A1	2	
(c)	$r = 1$ $\theta = \frac{1}{2}\pi, \frac{1}{3}\pi$	B1 B1B1	3	PI Deduct 1 mark if extra solutions
(d)		B2,1F	2	Positions of the 3 points relative to each other, must be approximately correct
(e)	$\text{Arg}(z_1 + z_2) = \frac{5}{12}\pi$ $\tan \frac{5}{12}\pi = \frac{1 + \frac{1}{2}\sqrt{3}}{\frac{1}{2}}$ $= 2 + \sqrt{3}$	B1 M1 A1	3	Clearly shown Allow if BO earned AG must earn BO for this
Total			12	

MFP2 (cont)

Q	Solution	Marks	Total	Comments
<p>4(a)</p> <p>Assume result true for $n = k$</p> $\sum_{r=1}^k (r+1)2^{r-1} = k2^k$ $\sum_{r=1}^{k+1} (r+1)2^{r-1} = k2^k + (k+2)2^k$ $= 2^k (k+k+2)$ $= 2^k (2k+2)$ $= 2^{k+1} (k+1)$ <p>$n = 1 \quad 2 \times 2^0 = 2 = 1 \times 2^1$</p> <p>$P_k \Rightarrow P_{k+1}$ and P_1 is true</p>	<p>M1A1</p> <p>m1</p> <p>A1</p> <p>B1</p> <p>E1</p>	<p>6</p>	<p>6</p>	<p>Provided previous 5 marks earned</p>
<p>(b)</p> $\sum_{r=1}^{2n} (r+1)2^{r-1} - \sum_{r=1}^n (r+1)2^{r-1}$ $= 2n \cdot 2^{2n} - n2^n$ $= n(2^{n+1} - 1)2^n$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>3</p>	<p>3</p>	<p>Sensible attempt at the difference between 2 series</p> <p>AG</p>
	Total		9	

MFP2 (cont)

Q	Solution	Marks	Total	Comments
5(a)		<p>B1</p> <p>B1</p> <p>B1</p>	<p>3</p>	<p>Circle</p> <p>Correct centre</p> <p>Touching both axes</p>
(b)	$ z _{\max} = OK$ $= \sqrt{4^2 + 4^2} + 4$ $= 4(\sqrt{2} + 1)$	<p>M1</p> <p>A1F</p> <p>A1F</p>	<p>3</p>	<p>Accept $\sqrt{4^2 + 4^2} + 4$ as a method</p> <p>Follow through circle in incorrect position</p> <p>AG</p>
(c)	<p>Correct position of z, ie L</p> $a = -\left(4 - 4\cos\frac{1}{6}\pi\right)$ $= -(4 - 2\sqrt{3})$ $b = 4 + 4\sin\frac{1}{6}\pi = 6$	<p>M1</p> <p>A1F</p> <p>A1F</p>	<p>3</p>	<p>Follow through circle in incorrect position</p>
	Total		9	

MFP2 (cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	$z + \frac{1}{z} = \cos \theta + i \sin \theta +$ $\cos(-\theta) + i \sin(-\theta)$	M1	2	AG
	$= 2 \cos \theta$	A1		
(ii)	$z^2 + \frac{1}{z^2} = \cos 2\theta + i \sin 2\theta$ $+ \cos(-2\theta) + i \sin(-2\theta)$	M1	2	OE
	$= 2 \cos 2\theta$	A1		
(iii)	$z^2 - z + 2 - \frac{1}{z} + \frac{1}{z^2}$ $= 2 \cos 2\theta - 2 \cos \theta + 2$	M1	3	AG
	Use of $\cos 2\theta = 2 \cos^2 \theta - 1$	m1		
	$= 4 \cos^2 \theta - 2 \cos \theta$	A1		
(b)	$z + \frac{1}{z} = 0 \quad z = \pm i$	M1A1	5	Alternative: $\cos \theta = 0 \quad \theta = \pm \frac{1}{2} \pi$ M1 $z = \pm i$ A1 $\cos \theta = \frac{1}{2} \quad \theta = \pm \frac{1}{3} \pi$ M1 $z = e^{\pm \frac{1}{3} \pi i} = \frac{1}{2} (1 \pm i\sqrt{3})$ A1 A1
	$z + \frac{1}{z} = 1 \quad z^2 - z + 1 = 0$	M1A1		
	$z = \frac{1 \pm i\sqrt{3}}{2}$	A1F		
	Accept solution to (b) if done otherwise			
	Alternative			
	If $\theta = + \frac{1}{2} \pi \quad \theta = - \frac{1}{2} \pi$	M1		
	$z = i \quad z = \frac{1 + \sqrt{3}i}{2}$	A1		
	Or any correct z values of θ	M1		
	Any 2 correct answers	A1		
	One correct answer only	B1		
Total			12	

MFP2 (cont)

Q	Solution	Marks	Total	Comments
7(a)(i)	$2\left(\frac{e^\theta - e^{-\theta}}{2}\right)\left(\frac{e^\theta + e^{-\theta}}{2}\right)$ $= \frac{e^{2\theta} - e^{-2\theta}}{2} = \sinh 2\theta$	M1A1	2	AG
(ii)	$\left(\frac{e^\theta - e^{-\theta}}{2}\right)^2 + \left(\frac{e^\theta + e^{-\theta}}{2}\right)^2$ $= \frac{e^{2\theta} - 2 + e^{-2\theta} + e^{2\theta} + 2 + e^{-2\theta}}{4}$ $= \cosh 2\theta$	M1 A1 A1	3	AG
(b)(i)	$\dot{x} = 3\cosh^2 \theta \sinh \theta''$ $\dot{y} = 3\sinh^2 \theta \cosh \theta$ $\dot{x}^2 + \dot{y}^2 = 9\cosh^4 \theta \sinh^2 \theta + 9\sinh^4 \theta \cosh^2 \theta$ $= 9\sinh^2 \theta \cosh^2 \theta (\cosh^2 \theta + \sinh^2 \theta)$ $= \frac{9}{4} \sinh^2 2\theta \cosh 2\theta$	M1A1 A1 M1 A1 A1	6	<p>Allow M1 for reasonable attempt at differentiation, but M0 for putting in terms of $e^{k\theta}$ or $\sinh 3\theta$ unless real progress made towards $\dot{x}^2 + \dot{y}^2$</p> <p>Allow this M1 if not squared out, must be clear sum in question is $\dot{x}^2 + \dot{y}^2$</p> <p>AG</p> <p>Accept $\int_0^1 \sqrt{\frac{9}{4} \sinh^2 2\theta \cosh 2\theta} d\theta$ but limits must appear somewhere</p> <p>AG</p>
(ii)	$S = \int_0^1 \frac{3}{2} \sinh 2\theta \sqrt{\cosh 2\theta} d\theta$ $u = \cosh 2\theta \quad du = 2 \sinh 2\theta d\theta$ $I = \int \frac{3}{4} u^{\frac{1}{2}} du = \frac{3}{4} \times \frac{2}{\frac{3}{2}} u^{\frac{3}{2}}$ $S = \left\{ \frac{1}{2} (\cosh 2\theta)^{\frac{3}{2}} \right\}_0^1$ $= \frac{1}{2} \left\{ (\cosh 2)^{\frac{3}{2}} - 1 \right\}$	M1 M1A1 A1F A1F A1	6	AG
	Total		17	
	TOTAL		75	