Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Level Examination January 2011

# **Mathematics**

MFP2

**Unit Further Pure 2** 

Wednesday 19 January 2011 1.30 pm to 3.00 pm

#### For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

#### Time allowed

• 1 hour 30 minutes

## Instructions


- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost
- Do all rough work in this book. Cross through any work that you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

#### **Advice**

 Unless stated otherwise, you may quote formulae, without proof, from the booklet.



# Answer all questions in the spaces provided.

1 (a) Sketch on an Argand diagram the locus of points satisfying the equation

$$|z-4+3i|=5$$
 (3 marks)

(b) (i) Indicate on your diagram the point P representing  $z_1$ , where both

$$|z_1 - 4 + 3i| = 5$$
 and  $\arg z_1 = 0$  (1 mark)

(ii) Find the value of  $|z_1|$ . (1 mark)

QUESTION PART REFERENCE	
•	
••••••	
•••••	
••••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



2 (a) Given that

$$u_r = \frac{1}{6}r(r+1)(4r+11)$$

show that

$$u_r - u_{r-1} = r(2r+3)$$
 (3 marks)

(b) Hence find the sum of the first hundred terms of the series

$$1 \times 5 + 2 \times 7 + 3 \times 9 + \dots + r(2r+3) + \dots$$
 (3 marks)

QUESTION PART REFERENCE	
•••••	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



- 3 (a) Show that  $(1+i)^3 = 2i 2$ . (2 marks)
  - **(b)** The cubic equation

$$z^3 - (5+i)z^2 + (9+4i)z + k(1+i) = 0$$

where k is a real constant, has roots  $\alpha$ ,  $\beta$  and  $\gamma$ .

It is given that  $\alpha = 1 + i$ .

- (i) Find the value of k. (3 marks)
- (ii) Show that  $\beta + \gamma = 4$ . (1 mark)
- (iii) Find the values of  $\beta$  and  $\gamma$ . (5 marks)

QUESTION PART REFERENCE	



QUESTION PART REFERENCE	
REFERENCE	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	



4 (a	Prove that the curve	
	$y = 12\cosh x - 8\sinh x - x$	
	has exactly one stationary point. (7 m	narks)
(b		= 9 . narks)
QUESTION PART REFERENCE		
•••••		
•••••		



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



- 5 (a) Given that  $u = \sqrt{1 x^2}$ , find  $\frac{du}{dx}$ . (2 marks)
  - **(b)** Use integration by parts to show that

$$\int_0^{\frac{\sqrt{3}}{2}} \sin^{-1} x \, dx = a\sqrt{3} \, \pi + b$$

where a and b are rational numbers.

(6 marks)

QUESTION PART REFERENCE	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



6 (a) Given that

$$x = \ln(\sec t + \tan t) - \sin t$$

show that

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \sin t \tan t \tag{4 marks}$$

**(b)** A curve is given parametrically by the equations

$$x = \ln(\sec t + \tan t) - \sin t$$
,  $y = \cos t$ 

The length of the arc of the curve between the points where t = 0 and  $t = \frac{\pi}{3}$  is denoted by s.

Show that  $s = \ln p$ , where p is an integer.

(6 marks)

QUESTION PART REFERENCE	
•••••	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	



7 (a) Given that

$$f(k) = 12^k + 2 \times 5^{k-1}$$

show that

$$f(k+1) - 5f(k) = a \times 12^k$$

where a is an integer.

(3 marks)

(b) Prove by induction that  $12^n + 2 \times 5^{n-1}$  is divisible by 7 for all integers  $n \ge 1$ .

QUESTION PART REFERENCE	
DADT	
FAILI	
REFERENCE	
• • • • • • • • • • • • • • • • • • • •	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	



- **8 (a)** Express in the form  $re^{i\theta}$ , where r > 0 and  $-\pi < \theta \le \pi$ :
  - (i)  $4(1+i\sqrt{3})$ ;

(ii) 
$$4(1-i\sqrt{3})$$
. (3 marks)

(b) The complex number z satisfies the equation

$$(z^3 - 4)^2 = -48$$

Show that 
$$z^3 = 4 \pm 4\sqrt{3}i$$
. (2 marks)

(c) (i) Solve the equation

$$(z^3-4)^2=-48$$

giving your answers in the form  $re^{i\theta}$ , where r > 0 and  $-\pi < \theta \le \pi$ . (5 marks)

- (ii) Illustrate the roots on an Argand diagram. (3 marks)
- (d) (i) Explain why the sum of the roots of the equation

$$(z^3 - 4)^2 = -48$$

is zero. (1 mark)

	Deduce that	π	$3\pi$	$5\pi$	$7\pi$	1	(2)
(11)		$\cos \frac{\pi}{9} + c$	$\cos \frac{\pi}{9} + c$	$\cos \frac{\pi}{9} + \epsilon$	$\cos \frac{\pi}{9} =$	$=\frac{1}{2}$ .	(3 marks



QUESTION PART REFERENCE	



QUESTION PART REFERENCE	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	
•••••	
••••••	



	END OF QUESTIONS
•••••	
•••••	
•••••	
QUESTION PART REFERENCE	
QUESTION	



