Centre Number			Candidate Number		
Surname					
Other Names					
Candidate Signature					



General Certificate of Education Advanced Level Examination January 2011

# **Mathematics**

MFP3

**Unit Further Pure 3** 

Monday 24 January 2011 9.00 am to 10.30 am

### For this paper you must have:

• the blue AQA booklet of formulae and statistical tables. You may use a graphics calculator.

#### Time allowed

• 1 hour 30 minutes

#### Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost
- Do all rough work in this book. Cross through any work that you do not want to be marked.

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

#### **Advice**

 Unless stated otherwise, you may quote formulae, without proof, from the booklet.

For Exam	iner's Use
Examine	r's Initials
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	



# Answer all questions in the spaces provided.

1 The function y(x) satisfies the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{f}(x, y)$$

where

$$f(x, y) = x + \sqrt{y}$$

and

$$y(3) = 4$$

Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where  $k_1 = hf(x_r, y_r)$  and  $k_2 = hf(x_r + h, y_r + k_1)$  and h = 0.1, to obtain an approximation to y(3.1), giving your answer to three decimal places. (5 marks)

QUESTION PART REFERENCE	
DADT	
PARI	
REFERENCE	
l	
<b></b>	



QUESTION PART REFERENCE	
••••••	
••••••	
•••••	
•••••	
•••••	



2 (a	Find the values of the constants $p$ and $q$ for which $p \sin x + q \cos x$ integral of the differential equation	is a particular
	$\frac{\mathrm{d}y}{\mathrm{d}x} + 5y = 13\cos x$	(3 marks)
(b	Hence find the general solution of this differential equation.	(3 marks)
QUESTION PART REFERENCE		
•••••		
•••••		
•••••		
		••••••



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



3	A curve C has polar equation $r(1 + \cos \theta) = 2$ .
(a	Find the cartesian equation of C, giving your answer in the form $y^2 = f(x)$ .  (5 marks)
(b	The straight line with polar equation $4r = 3 \sec \theta$ intersects the curve $C$ at the points $P$ and $Q$ . Find the length of $PQ$ . (4 marks)
QUESTION PART REFERENCE	
·······	
•	
••••••	
••••••	
•••••	
••••••	
••••••	
••••••	
•••••	



QUESTION PART REFERENCE	
REFERENCE	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	



			_					_
4	By using a	n integrating	factor,	find the	solution	of the	differential	equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} - \frac{2}{x}y = 2x^3 \mathrm{e}^{2x}$$

given that  $y = e^4$  when x = 2. Give your answer in the form y = f(x). (9 marks)

QUESTION PART	
DADT	
PEEEDENOE	
REFERENCE	
	• • • • • • • • • • • • • • • • • • • •
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • •	
• • • • • • • • • •	
• • • • • • • • • • •	
•••••	
•••••	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



- 5 (a) Write  $\frac{4}{4x+1} \frac{3}{3x+2}$  in the form  $\frac{C}{(4x+1)(3x+2)}$ , where C is a constant.
  - **(b)** Evaluate the improper integral

$$\int_{1}^{\infty} \frac{10}{(4x+1)(3x+2)} \, \mathrm{d}x$$

showing the limiting process used and giving your answer in the form  $\ln k$ , where k is a constant. (6 marks)


QUESTION PART REFERENCE	
PART	
REFERENCE	
• • • • • • • • • • • • • • • • • • • •	
,	



QUESTION PART REFERENCE	
••••••	
•••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



6 The diagram shows a sketch of a curve C.



The polar equation of the curve is

$$r = 2\sin 2\theta \sqrt{\cos \theta}$$
,  $0 \le \theta \le \frac{\pi}{2}$ 

Show that the area of the region bounded by C is  $\frac{16}{15}$ .

(7 marks)

QUESTION PART REFERENCE	
PART	
NLI LINLINGL	
• • • • • • • • • • • • • • • • • • • •	



QUESTION PART REFERENCE	
•	
•••••	
••••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



7 (a) Write down the expansions in ascending powers of x up to and including the term in  $x^3$  of:

(i) 
$$\cos x + \sin x$$
; (1 mark)

(ii) 
$$\ln(1+3x)$$
. (1 mark)

**(b)** It is given that  $y = e^{\tan x}$ .

(i) Find 
$$\frac{dy}{dx}$$
 and show that  $\frac{d^2y}{dx^2} = (1 + \tan x)^2 \frac{dy}{dx}$ . (5 marks)

(ii) Find the value of 
$$\frac{d^3y}{dx^3}$$
 when  $x = 0$ . (2 marks)

(iii) Hence, by using Maclaurin's theorem, show that the first four terms in the expansion, in ascending powers of x, of  $e^{\tan x}$  are

$$1 + x + \frac{1}{2}x^2 + \frac{1}{2}x^3$$
 (2 marks)

(c) Find

$$\lim_{x \to 0} \left[ \frac{e^{\tan x} - (\cos x + \sin x)}{x \ln(1 + 3x)} \right]$$
 (3 marks)

QUESTION	
QUESTION PART	
REFERENCE	



QUESTION	
QUESTION PART REFERENCE	
KEFEKENCE	
	l l
• • • • • • • • • • • • • • • • • • • •	
	I
	l l
• • • • • • • • • • • • • • • • • • • •	
•••••	
	I
	l la companya di managantan di managantan di managantan di managantan di managantan di managantan di managanta
• • • • • • • • • • • • • • • • • • • •	
	l l
	l la companya di managanta di ma
•••••	
	l l
	l l
• • • • • • • • • • • • • • • • • • • •	



QUESTION PART REFERENCE	
••••••	
••••••	
••••••	
••••••	
••••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	
•••••	



QUESTION PART REFERENCE	



8 (a) Given that  $x = e^t$  and that y is a function of x, show that

$$x\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}y}{\mathrm{d}t} \tag{2 marks}$$

(b) Hence show that the substitution  $x = e^t$  transforms the differential equation

$$x^2 \frac{\mathrm{d}^2 y}{\mathrm{d}x^2} - 3x \frac{\mathrm{d}y}{\mathrm{d}x} + 4y = 2\ln x$$

into

$$\frac{\mathrm{d}^2 y}{\mathrm{d}t^2} - 4\frac{\mathrm{d}y}{\mathrm{d}t} + 4y = 2t \tag{5 marks}$$

(c) Find the general solution of the differential equation

$$\frac{\mathrm{d}^2 y}{\mathrm{d}t^2} - 4\frac{\mathrm{d}y}{\mathrm{d}t} + 4y = 2t \tag{6 marks}$$

(d) Hence solve the differential equation  $x^2 \frac{d^2y}{dx^2} - 3x \frac{dy}{dx} + 4y = 2 \ln x$ , given that  $y = \frac{3}{2}$  and  $\frac{dy}{dx} = \frac{1}{2}$  when x = 1. (5 marks)

PART REFERENCE	



QUESTION PART REFERENCE	
•••••	
•••••	
•••••	
•••••	
•••••	



QUESTION PART	
REFERENCE	
•••••	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
• • • • • • • • • • • • • • • • • • • •	
	END OF QUESTIONS
Copyrig	ht © 2011 AQA and its licensors. All rights reserved.

