

Centre Number						Candidate Number				
Surname										
Other Names										
Candidate Signature										

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
TOTAL	



General Certificate of Education
Advanced Level Examination
June 2011

Mathematics

MFP3

Unit Further Pure 3

Thursday 16 June 2011 1.30 pm to 3.00 pm

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed

- 1 hour 30 minutes

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.



J U N 1 1 M F P 3 0 1

Answer **all** questions in the spaces provided.

1 The function $y(x)$ satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where $f(x, y) = x + \ln(1 + y)$

and $y(2) = 1$

Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where $k_1 = hf(x_r, y_r)$ and $k_2 = hf(x_r + h, y_r + k_1)$ and $h = 0.2$, to obtain an approximation to $y(2.2)$, giving your answer to four decimal places. (5 marks)

QUESTION
PART
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- 2 (a)** Find the values of the constants p and q for which $p + qxe^{-2x}$ is a particular integral of the differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 2y = 4 - 9e^{-2x} \quad (5 \text{ marks})$$

- (b)** Hence find the general solution of this differential equation. (3 marks)

- (c)** Hence express y in terms of x , given that $y = 4$ when $x = 0$ and that $\frac{dy}{dx} \rightarrow 0$ as $x \rightarrow \infty$. (4 marks)

QUESTION
PART
REFERENCE



QUESTION
PART
REFERENCE

This section contains a large rectangular area with a solid left border and a solid bottom border. The interior is filled with horizontal dotted lines, providing space for writing answers to questions.

Turn over ►



3 (a) Find $\int x^2 \ln x \, dx$. *(3 marks)*

(b) Explain why $\int_0^e x^2 \ln x \, dx$ is an improper integral. *(1 mark)*

(c) Evaluate $\int_0^e x^2 \ln x \, dx$, showing the limiting process used. *(3 marks)*

QUESTION
PART
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



4 By using an integrating factor, find the solution of the differential equation

$$\frac{dy}{dx} + (\cot x)y = \sin 2x, \quad 0 < x < \frac{\pi}{2}$$

given that $y = \frac{1}{2}$ when $x = \frac{\pi}{6}$.

(10 marks)

QUESTION
PART
REFERENCE



5 (a) Given that $y = \ln(1 + 2 \tan x)$, find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.

(You may leave your expression for $\frac{d^2y}{dx^2}$ unsimplified.) (4 marks)

(b) Hence, using Maclaurin's theorem, find the first two non-zero terms in the expansion, in ascending powers of x , of $\ln(1 + 2 \tan x)$. (2 marks)

(c) Find

$$\lim_{x \rightarrow 0} \left[\frac{\ln(1 + 2 \tan x)}{\ln(1 - x)} \right] \quad (4 \text{ marks})$$

QUESTION
PART
REFERENCE



6 A differential equation is given by

$$(x^3 + 1) \frac{d^2y}{dx^2} - 3x^2 \frac{dy}{dx} = 2 - 4x^3$$

(a) Show that the substitution

$$u = \frac{dy}{dx} - 2x$$

transforms this differential equation into

$$(x^3 + 1) \frac{du}{dx} = 3x^2 u \qquad (4 \text{ marks})$$

(b) Hence find the general solution of the differential equation

$$(x^3 + 1) \frac{d^2y}{dx^2} - 3x^2 \frac{dy}{dx} = 2 - 4x^3$$

giving your answer in the form $y = f(x)$. (8 marks)

QUESTION
PART
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



7 The curve C_1 is defined by $r = 2 \sin \theta$, $0 \leq \theta < \frac{\pi}{2}$.

The curve C_2 is defined by $r = \tan \theta$, $0 \leq \theta < \frac{\pi}{2}$.

(a) Find a cartesian equation of C_1 . (3 marks)

(b) (i) Prove that the curves C_1 and C_2 meet at the pole O and at one other point, P , in the given domain. State the polar coordinates of P . (4 marks)

(ii) The point A is the point on the curve C_1 at which $\theta = \frac{\pi}{4}$.

The point B is the point on the curve C_2 at which $\theta = \frac{\pi}{4}$.

Determine which of the points A or B is further away from the pole O , justifying your answer. (2 marks)

(iii) Show that the area of the region bounded by the arc OP of C_1 and the arc OP of C_2 is $a\pi + b\sqrt{3}$, where a and b are rational numbers. (10 marks)

QUESTION
PART
REFERENCE



QUESTION
PART
REFERENCE

Area with horizontal dotted lines for writing.

END OF QUESTIONS

Copyright © 2011 AQA and its licensors. All rights reserved.

