



**General Certificate of Education (A-level)
June 2012**

Mathematics

MFP4

(Specification 6360)

Further Pure 4

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2012 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Q	Solution	Marks	Total	Comments
1	<p>Attempt at $\begin{vmatrix} 3 & 2 & p \\ 7 & -1 & 6 \\ 2 & 1 & 3 \end{vmatrix} = 0$</p> <p>Solving a linear equation in p $p = 5$</p> <p>ALT Lin.dep. iff \exists constants a and b s.t. $a\mathbf{u} + b\mathbf{v} = \mathbf{w}$</p> <p>Solving simultaneously $3a + 7b = 2$ and $2a - b = 1$ M1 $a = \frac{9}{17}$, $b = \frac{1}{17}$ A1</p> <p>Substituting back into \mathbf{k} component ($ap + 6b = 3$) to find “their” p correctly A1F</p>	M1 M1 A1	3	(from \mathbf{i} and \mathbf{j} components)
Total			3	
2(a)	<p>Choice of $\begin{pmatrix} 4 \\ 7 \\ -4 \end{pmatrix}$ as direction vector</p> <p>$\sqrt{4^2 + 7^2 + 4^2}$ or $\sqrt{3^2 + 2^2 + 6^2}$ $\frac{4}{9}$, $\frac{7}{9}$, $-\frac{4}{9}$ or $\frac{3}{7}$, $-\frac{2}{7}$, $\frac{6}{7}$</p>	B1 M1 A1	3	Either attempted ft their chosen direction vector
(b)	Direction cosines are the cosines of the angles between the line and the coordinate axes	B1	1	
Total			4	
3(a)	<p>eg $\begin{vmatrix} yz & x(y+z) & xy \\ x & y+z & z \\ x^2 & z^2 - y^2 & z^2 \end{vmatrix}$</p> <p>$= (y+z) \begin{vmatrix} yz & x & xy \\ x & 1 & z \\ x^2 & z-y & z^2 \end{vmatrix}$</p>	M1 A1	2	$C_2' = C_2 + C_3$
(b)	<p>eg $(y+z) \begin{vmatrix} y(z-x) & x & xy \\ x-z & 1 & z \\ x^2 - z^2 & z-y & z^2 \end{vmatrix}$</p> <p>$C_1' = C_1 - C_3$</p> <p>$= (x-z)(y+z) \begin{vmatrix} -y & x & xy \\ 1 & 1 & z \\ x+z & z-y & z^2 \end{vmatrix}$</p> <p>$= (x-z)(y+z) \begin{vmatrix} -(x+y) & x & xy \\ 0 & 1 & z \\ x+y & z-y & z^2 \end{vmatrix}$</p> <p>$= (x-z)(y+z)(x+y)(xz - xy - yz)$</p>	M1 A1 M1 A1	4	Attempt at a second linear factor $C_1' = C_1 - C_2$ Complete attempt at remaining factors (M0 if they just expand and can do nothing with it)
Total			6	

Q	Solution	Marks	Total	Comments
4(a)	$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -4 & 7 \\ 2 & -2 & 3 \end{vmatrix} = \begin{bmatrix} 2 \\ 5 \\ 2 \end{bmatrix}$	M1		Attempt at the vector product of the 2 d.v.s OR two scalar products = 0 and some manipulation attempt
		A1	2	
(b)(i)	$\overrightarrow{AB} = \begin{bmatrix} 2\beta + 7 \\ 19 - 2\beta \\ 3\beta - 2 \end{bmatrix} - \begin{bmatrix} 3\alpha + 7 \\ -25 - 4\alpha \\ 7\alpha + 9 \end{bmatrix}$ $= \begin{bmatrix} 2\beta - 3\alpha \\ 44 - 2\beta + 4\alpha \\ -11 + 3\beta - 7\alpha \end{bmatrix}$	M1		Good attempt
	$\overrightarrow{AB} = \lambda \mathbf{n} = \begin{bmatrix} 2\lambda \\ 5\lambda \\ 2\lambda \end{bmatrix}$ <p>Legitimately getting given system of equations:</p> $3\alpha - 2\beta + 2\lambda = 0$ $4\alpha - 2\beta - 5\lambda = -44$ $7\alpha - 3\beta + 2\lambda = -11$	M1		With their \overrightarrow{AB} (involving α and β) and their \mathbf{n}
(ii)	Solving this 3×3 system in α , β and λ (possibly just λ given or both α , β)	A1	3	
	For $\alpha = -2$, $\beta = 3$ ($\lambda = 6$)	A1		
	$A = (1, -17, -5)$ and $B = (13, 13, 7)$	A1	3	Give one A1 for a correct pair (α , A) or (β , B)
(iii)	Shortest distance = $\sqrt{12^2 + 30^2 + 12^2}$ or $ \lambda \mathbf{n} = 6\sqrt{2^2 + 5^2 + 2^2}$	M1		SC: allow all 3 marks ft for misreads of the signs (44 and/or 11 only) Allow also for the shortest distance formula $ (\mathbf{b} - \mathbf{a}) \cdot \hat{\mathbf{n}} $
	$6\sqrt{33}$, $\frac{198}{\sqrt{33}}$, $\sqrt{1188}$ or AWRT 34.5	A1	2	CAO
	Total		10	

Q	Solution	Marks	Total	Comments
5(a)(i)	Char. eqn. is $\lambda^2 - 2\lambda + 1 = 0$	M1		Written down or attempted via determinant
	$\lambda = 1$ (twice)	A1		
	Substituting $\lambda = 1 \Rightarrow -12x + 9y = 0$ and/or $-16x + 12y = 0$	M1		
	Eigenvector(s) $\begin{bmatrix} 3 \\ 4 \end{bmatrix}$	A1	4	Any non-zero multiple will do
(ii)	$m = \frac{4}{3}$	B1		ft
	LOIPs since $\lambda = 1$ (or full description)	B1	2	
(iii)	$\begin{bmatrix} -11 & 9 \\ -16 & 13 \end{bmatrix} \begin{bmatrix} x \\ \frac{4}{3}x + c \end{bmatrix}$	M1		m must be their numerical value
	$= \begin{bmatrix} -11x + 9(\frac{4}{3}x + c) \\ -16x + 13(\frac{4}{3}x + c) \end{bmatrix}$ or $\begin{bmatrix} x + 9c \\ \frac{4}{3}x + 13c \end{bmatrix}$	A1		ft
	For showing $y' = \frac{4}{3}x' + c$	B1	3	Impossible without correct prior working
(b)	... parallel to $y = \frac{4}{3}x$ "	B1		ft their m
	For mapping any one point to a correct image point, eg (1, 0) to (-11, -16), (0, 1) to (9, 13) or (1, 1) to (-2, -3)	B1	2	If they cite multiple points, including incorrect ones, then B0
(c)	Shear parallel to $y = \frac{4}{3}x$	M1		MUST be the same LOIPs as in (b)
	For mapping any one point to a correct image point	A1	2	ft incorrect ones ONLY if they are the reverse of those cited in (b)
Total			13	

Q	Solution	Marks	Total	Comments
6(a)	For attempt at/getting normal d.v. $\begin{bmatrix} 1 \\ 5 \\ 7 \end{bmatrix}$	M1 A1	5	Condone lack of r or r here
	For getting a point on the line eg (0, -8, -11)	M1 A1		
	For line equation: (r =) (their point) + λ (their d.v.)	B1		
	ALT eg Adding I_1 and I_2 M1 $\Rightarrow 5x - y = 8$ A1 Setting (eg) $x = \lambda$ and getting y, z in terms of λ : $y = 5\lambda - 8, z = 7\lambda - 11$ M1 Turning this into a vector equation of the given form M1			
	(r =) $\begin{bmatrix} 0 \\ -8 \\ -11 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 5 \\ 7 \end{bmatrix}$ A1			
(b)(i)	Substituting their x, y, z in terms of λ into I_3 's equation ($12x - y - z = 40$)	M1		
	For correct statement with no λ 's in: ($19 = 40$)	A1		ft on incorrect values from their "point"
	Correct conclusion, from valid working, that the system is inconsistent	B1	3	or "consistent" if it genuinely yields $40 = 40$
(ii)	The three planes form a (triangular) prism – allow clear diagram	B1	1	ft "sheaf" from a "consistent" conclusion
(c)(i)	e.g. $I_2 + I_3 \Rightarrow 15x - 3y = 45$ Setting $x = 0$ (eg)	M1 M1		
	Any correct point, eg (0, -15, -25), (3, 0, -4), $(\frac{25}{7}, \frac{20}{7}, 0)$, (4, 5, 3) etc	A1	3	
(ii)	r = $\begin{bmatrix} 3 \\ 0 \\ -4 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 5 \\ 7 \end{bmatrix}$	B1	1	ft their common point and d.v. from (a) Penalise missing r or r here
			13	

Q	Solution	Marks	Total	Comments
7(a)(i)	$\mathbf{A}^T = \begin{bmatrix} k & 2 & 1 \\ 1 & k & 2 \\ 2 & 1 & k \end{bmatrix}$ <p>Good multiplication attempt at $\mathbf{A} \mathbf{A}^T$</p> $= \begin{bmatrix} k^2 + 5 & 3k + 2 & 3k + 2 \\ 3k + 2 & k^2 + 5 & 3k + 2 \\ 3k + 2 & 3k + 2 & k^2 + 5 \end{bmatrix}$ <p>$k = -\frac{2}{3}$ $m = 5\frac{4}{9}$ or $\frac{49}{9}$</p>	B1 M1 A1 A1 A1A1	6	Main diagonal correct All others correct If they multiply $\mathbf{A}^T \mathbf{A}$ instead, they can score B1 M1 A0 A0 A1 A1
(ii)	$\begin{bmatrix} -\frac{2}{3} & 1 & 2 \\ 2 & -\frac{2}{3} & 1 \\ 1 & 2 & -\frac{2}{3} \end{bmatrix}^{-1} = \frac{9}{49} \begin{bmatrix} -\frac{2}{3} & 2 & 1 \\ 1 & -\frac{2}{3} & 2 \\ 2 & 1 & -\frac{2}{3} \end{bmatrix}$ <p>Accept $\frac{9}{49} \begin{bmatrix} k & 2 & 1 \\ 1 & k & 2 \\ 2 & 1 & k \end{bmatrix}$ since the value of k is now known, but not just $\frac{9}{49} \mathbf{A}^T$</p> <p>Decimal version (correct to at least 3sf) is also ok:</p> $\begin{bmatrix} -0.122 & 0.367 & 0.184 \\ 0.184 & -0.122 & 0.367 \\ 0.367 & 0.184 & -0.122 \end{bmatrix}$	B1	1	ft $\frac{1}{\text{their } m}$ and their k
(b)(i)	$\det \mathbf{A} = k^3 - 6k + 9$	M1 A1	2	Good attempt (cubic)
(ii)	$\det \mathbf{A} = (k + 3)(k^2 - 3k + 3)$ $k = -3$ $\Delta = 9 - 12 < 0 \Rightarrow$ no further real roots	M1 A1 B1	3	Factorisation attempt Special Cases $k = -3$ but no working B1 For all 3 roots, -3 and $\frac{3+i\sqrt{3}}{2}$ given with no supporting working B3
(iii)	<p>Replacing k by $(k - 7)$</p> <p>$k = 4$ (however obtained)</p>	M1 A1	2	NOT just in the determinant form (ie starting again) ft their previous $k + 7$
			14	

Q	Solution	Marks	Total	Comments
8(a)	$\begin{bmatrix} 7 \\ 4 \\ 6 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = 14 + 4 + 18 = 36 \Rightarrow Q \text{ in } \Pi$	B1	1	
(b)	$\begin{bmatrix} -7 \\ 5 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = -14 + 5 + 9 = 0$ <p>Explanation that l is perpendicular to Π's normal $\Rightarrow l$ is parallel to Π</p>	B1 B1	2	Shown
	<p>ALT</p> $\begin{bmatrix} 20 - 7\mu \\ 5\mu - 8 \\ 3\mu + 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = 35 + 0\mu \quad \text{B1}$ <p>Explanation that, since this $\neq 36$ (and constant), l does not intersect Π and must therefore be parallel to it B1</p>			
(c)	Mark according to whichever scheme gives the greatest credit			
	MARK SCHEME I			
	<p>For $\mathbf{p} = \mathbf{q} + \lambda \mathbf{n} = \begin{bmatrix} 7 + 2\lambda \\ 4 + \lambda \\ 6 + 3\lambda \end{bmatrix}$ for some λ</p>	B1		
	<p>For $PQ = \lambda \sqrt{14}$ or equivalent</p>	B1		
	$\overline{PR} = \mathbf{r} - \mathbf{p} = \begin{bmatrix} 20 - 7\mu \\ 5\mu - 8 \\ 3\mu + 1 \end{bmatrix} - \begin{bmatrix} 7 + 2\lambda \\ 4 + \lambda \\ 6 + 3\lambda \end{bmatrix}$	M1		
	$= \begin{bmatrix} -7\mu - 2\lambda + 13 \\ 5\mu - \lambda - 12 \\ 3\mu - 3\lambda - 5 \end{bmatrix}$	A1		
	<p>Setting $\overline{PR} \cdot \begin{bmatrix} -7 \\ 5 \\ 3 \end{bmatrix} = 0$</p>	M1		
	<p>Solving linear equation for λ</p>	M1		$\Rightarrow 83\mu - 166 = 0$ ultimately
	$\mu = 2 \text{ and/or } \overline{PR} = \begin{bmatrix} -(2\lambda + 1) \\ -(\lambda + 2) \\ 1 - 3\lambda \end{bmatrix}$	A1		
	$PR^2 = PQ^2$	M1		$(1 + 2\lambda)^2 + (2 + \lambda)^2 + (1 - 3\lambda)^2 = 14\lambda^2$
	$(\lambda = -3) \quad P = (1, 1, -3)$	A1	9	

Q	Solution	Marks	Total	Comments
8(c) cont	<p>MARK SCHEME II</p> <p>For $\mathbf{p} = \mathbf{q} + \lambda \mathbf{n} = \begin{bmatrix} 7+2\lambda \\ 4+\lambda \\ 6+3\lambda \end{bmatrix}$ for some λ</p> <p>For $PQ^2 = 14\lambda^2$</p> <p>$\overline{PR} = \mathbf{r} - \mathbf{p} = \begin{bmatrix} 20-7\mu \\ 5\mu-8 \\ 3\mu+1 \end{bmatrix} - \begin{bmatrix} 7+2\lambda \\ 4+\lambda \\ 6+3\lambda \end{bmatrix}$</p> <p>$= \begin{bmatrix} -7\mu-2\lambda+13 \\ 5\mu-\lambda-12 \\ 3\mu-3\lambda-5 \end{bmatrix}$</p> <p>$PR^2 = (-13+7\mu+2\lambda)^2 + (12-5\mu+\lambda)^2 + (5-3\mu+3\lambda)^2$</p> <p>Setting $PR^2 = PQ^2$ $\Rightarrow 83\mu^2 - 332\mu + 338 + 2\lambda = 0$</p> <p>Considering discriminant = 0 or $83(\mu-2)^2 = -2(\lambda+3)$</p> <p>$\mu = 2, \lambda = -3$ and $P = (1, 1, -3)$</p> <p>MARK SCHEME III</p> <p>For $\mathbf{p} = \mathbf{q} + \lambda \mathbf{n} = \begin{bmatrix} 7+2\lambda \\ 4+\lambda \\ 6+3\lambda \end{bmatrix}$ for some λ</p> <p>For $PQ^2 = 14\lambda^2$</p> <p>For $\mathbf{r} = \begin{bmatrix} 20-7\mu \\ 5\mu-8 \\ 3\mu+1 \end{bmatrix}$</p> <p>For $QR^2 =$ $(13-7\mu)^2 + (5\mu-12)^2 + (3\mu-5)^2$ $= 83\mu^2 - 332\mu + 338 = 83(\mu-2)^2 + 6$</p> <p>For R closest to Q when $\mu = 2,$ $R = (6, 2, 7)$</p> <p>Then $\overline{RP} = \begin{bmatrix} 1+2\lambda \\ 2+\lambda \\ 3\lambda-1 \end{bmatrix}$</p> <p>Setting $PR^2 = PQ^2$ $\Rightarrow (1+2\lambda)^2 + (2+\lambda)^2 + (3\lambda-1)^2 = 14\lambda^2$</p> <p>$(\lambda = -3) \quad P = (1, 1, -3)$</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p>	<p>(9)</p> <p>(9)</p>	<p>Correct quadratic in μ</p> <p>ft</p>
	Total		12	
	TOTAL		75	