

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS**  
**Advanced Subsidiary General Certificate of Education**  
**Advanced General Certificate of Education**

**MATHEMATICS**  
Core Mathematics 4

**4724**

Monday      **23 JANUARY 2006**      Afternoon      1 hour 30 minutes

Additional materials:  
8 page answer booklet  
Graph paper  
List of Formulae (MF1)

**TIME**      1 hour 30 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

---

**This question paper consists of 3 printed pages and 1 blank page.**

- 1 Simplify  $\frac{x^3 - 3x^2}{x^2 - 9}$ . [3]
- 2 Given that  $\sin y = xy + x^2$ , find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ . [5]
- 3 (i) Find the quotient and the remainder when  $3x^3 - 2x^2 + x + 7$  is divided by  $x^2 - 2x + 5$ . [4]  
 (ii) Hence, or otherwise, determine the values of the constants  $a$  and  $b$  such that, when  $3x^3 - 2x^2 + ax + b$  is divided by  $x^2 - 2x + 5$ , there is no remainder. [2]
- 4 (i) Use integration by parts to find  $\int x \sec^2 x \, dx$ . [4]  
 (ii) Hence find  $\int x \tan^2 x \, dx$ . [3]
- 5 A curve is given parametrically by the equations  $x = t^2$ ,  $y = 2t$ .  
 (i) Find  $\frac{dy}{dx}$  in terms of  $t$ , giving your answer in its simplest form. [2]  
 (ii) Show that the equation of the tangent to the curve at  $(p^2, 2p)$  is  

$$py = x + p^2. \quad [2]$$
  
 (iii) Find the coordinates of the point where the tangent at  $(9, 6)$  meets the tangent at  $(25, -10)$ . [4]
- 6 (i) Show that the substitution  $x = \sin^2 \theta$  transforms  $\int \sqrt{\frac{x}{1-x}} \, dx$  to  $\int 2 \sin^2 \theta \, d\theta$ . [4]  
 (ii) Hence find  $\int_0^1 \sqrt{\frac{x}{1-x}} \, dx$ . [5]
- 7 The expression  $\frac{11 + 8x}{(2-x)(1+x)^2}$  is denoted by  $f(x)$ .  
 (i) Express  $f(x)$  in the form  $\frac{A}{2-x} + \frac{B}{1+x} + \frac{C}{(1+x)^2}$ , where  $A$ ,  $B$  and  $C$  are constants. [5]  
 (ii) Given that  $|x| < 1$ , find the first 3 terms in the expansion of  $f(x)$  in ascending powers of  $x$ . [5]

- 8 (i) Solve the differential equation

$$\frac{dy}{dx} = \frac{2-x}{y-3},$$

giving the particular solution that satisfies the condition  $y = 4$  when  $x = 5$ . [5]

- (ii) Show that this particular solution can be expressed in the form

$$(x-a)^2 + (y-b)^2 = k,$$

where the values of the constants  $a$ ,  $b$  and  $k$  are to be stated. [3]

- (iii) Hence sketch the graph of the particular solution, indicating clearly its main features. [3]

- 9 Two lines have vector equations

$$\mathbf{r} = \begin{pmatrix} 4 \\ 2 \\ -6 \end{pmatrix} + t \begin{pmatrix} -8 \\ 1 \\ -2 \end{pmatrix} \quad \text{and} \quad \mathbf{r} = \begin{pmatrix} -2 \\ a \\ -2 \end{pmatrix} + s \begin{pmatrix} -9 \\ 2 \\ -5 \end{pmatrix},$$

where  $a$  is a constant.

- (i) Calculate the acute angle between the lines. [5]

- (ii) Given that these two lines intersect, find  $a$  and the point of intersection. [8]