



General Certificate of Education

Mathematics 6360

MD02 Decision 2

Mark Scheme

2010 examination - January series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2010 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.



Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MD02

Q	Solution	Marks	Total	Comments																																												
1	 <p>Activity Node Data:</p> <table border="1"> <tr><th>Activity</th><th>ES</th><th>D</th><th>EF</th></tr> <tr><td>A</td><td>0</td><td>4</td><td>4</td></tr> <tr><td>B</td><td>0</td><td>2</td><td>2</td></tr> <tr><td>C</td><td>0</td><td>5</td><td>5</td></tr> <tr><td>D</td><td>4</td><td>3</td><td>7</td></tr> <tr><td>E</td><td>4</td><td>5</td><td>9</td></tr> <tr><td>F</td><td>5</td><td>4</td><td>9</td></tr> <tr><td>G</td><td>5</td><td>6</td><td>11</td></tr> <tr><td>H</td><td>9</td><td>6</td><td>15</td></tr> <tr><td>I</td><td>11</td><td>1</td><td>12</td></tr> <tr><td>J</td><td>15</td><td>2</td><td>17</td></tr> </table>	Activity	ES	D	EF	A	0	4	4	B	0	2	2	C	0	5	5	D	4	3	7	E	4	5	9	F	5	4	9	G	5	6	11	H	9	6	15	I	11	1	12	J	15	2	17			
Activity	ES	D	EF																																													
A	0	4	4																																													
B	0	2	2																																													
C	0	5	5																																													
D	4	3	7																																													
E	4	5	9																																													
F	5	4	9																																													
G	5	6	11																																													
H	9	6	15																																													
I	11	1	12																																													
J	15	2	17																																													
(a)(i)	Earliest start times	M1 A1	2	At least 4 correct (other than zeros) All correct																																												
(ii)	Latest finish times	M1 A1	2	Up to one error + follow through All correct																																												
(b)	Float for G = 3 days	B1✓	1	ft their G values																																												
(c)	Critical paths <i>A E H J</i> and <i>C F H J</i> Minimum completion time = 17 days	B1 B1 B1	3	First correct path Second path (and no others) Must be stated explicitly																																												
(d)	 <p>Activity Scheduling Data:</p> <table border="1"> <tr><th>Activity</th><th>Start</th><th>End</th><th>Workers</th></tr> <tr><td>A</td><td>0</td><td>4</td><td>2</td></tr> <tr><td>B</td><td>2</td><td>4</td><td>7</td></tr> <tr><td>C</td><td>0</td><td>5</td><td>4</td></tr> <tr><td>D</td><td>6</td><td>9</td><td>7</td></tr> <tr><td>E</td><td>4</td><td>9</td><td>3</td></tr> <tr><td>F</td><td>5</td><td>9</td><td>4</td></tr> <tr><td>G</td><td>8</td><td>14</td><td>8</td></tr> <tr><td>H</td><td>9</td><td>15</td><td>3</td></tr> <tr><td>I</td><td>14</td><td>15</td><td>8</td></tr> <tr><td>J</td><td>15</td><td>17</td><td>2</td></tr> </table>	Activity	Start	End	Workers	A	0	4	2	B	2	4	7	C	0	5	4	D	6	9	7	E	4	9	3	F	5	9	4	G	8	14	8	H	9	15	3	I	14	15	8	J	15	17	2			<p>One of their CPs – correct heights <i>A E H J</i> and <i>C F (H J)</i> correct <i>B</i> starting at 2 (and ending at 4) <i>D</i> starting at 6 (and ending at 9) <i>G</i> starting at 8 (and ending at 14) <i>I</i> starting at 14 (and ending at 15)</p> <p>M1 A1 M1 A1 A1</p> <p>5</p> <p>NB “holes” penalise first A1 earned One correct with correct height Two correct with correct height All correct with correct height and no slack Withhold first A1 earned if it is not clear which activities take place at any given time</p>
Activity	Start	End	Workers																																													
A	0	4	2																																													
B	2	4	7																																													
C	0	5	4																																													
D	6	9	7																																													
E	4	9	3																																													
F	5	9	4																																													
G	8	14	8																																													
H	9	15	3																																													
I	14	15	8																																													
J	15	17	2																																													
	Total		13																																													

MD02 (cont)

Q	Solution	Marks	Total	Comments
2(a)	8 7 9 10 8	B1	1	Adding extra row equal values
	9 x 8 7 11			
	12 10 9 9 10			
	11 9 8 11 11			
	12 12 12 12 12			
(b)(i)	0 0 1 3 0	B1✓	5	Reducing columns first
	1 $x-7$ 0 0 3			
	4 3 1 2 2			
	3 2 0 4 3			
	4 5 4 5 4			
	0 0 1 3 0	B1✓		Reducing rows
	1 $x-7$ 0 0 3			
	3 2 0 1 1			
	3 2 0 4 3			
	0 1 0 1 0			
Zeros covered with 4 lines (stated or drawn)		E1		or
0 0 1 3 0 1 $x-7$ 0 0 3 3 2 0 1 1 3 2 0 4 3 0 1 0 1 0	0 0 1 3 0 1 $x-7$ 0 0 3 3 2 0 1 1 3 2 0 4 3 0 1 0 1 0			
		M1		Augmentation – adding 1 to double covered and subtracting 1 from uncovered
⇒	0 0 2 3 0	A1	5	or
	1 $x-7$ 1 0 3			
	2 1 0 0 0			
	2 1 0 3 2			
	0 1 1 1 0			
(ii)	S1, V2, Z3, T4	M1		At least 2 matched correctly or “rings” on final tableau (Ron not assigned)
		A1	2	
(iii)	Total time 32 (minutes)	B1	1	
(c)	V3, T4, R1 or V3, T4, Z1	B1		First matching
		B1	2	Second matching and no other
Total			11	

MD02 (cont)

Q	Solution	Marks	Total	Comments
3(a)	Row min -2 -3 -5			Row minima and column maxima (all values)
	Col max 4 0 -2	B1		
	Max (row min) = -2 Min (col max) = -2	M1		Both attempted or stated/indicated
	Since these are equal, there is a stable solution	A1		Must have both values = -2 plus statement (withhold if max (min) and min (max) not stated)
	Ann plays A_1 and Bill plays B_3 for play-safe	E1	4	
	(b)(i) Let Russ play R_1 with probability p			And R_2 with probability $1-p$
	C_1 : expected gain $-4p + 2(1-p)$			$(2-6p)$
	C_2 : $7p - (1-p) = 8p-1$	M1		2 correct unsimplified
	C_3 : $3p + (1-p) = 1-4p$	A1		All correct
		M1 A1		Plotting 3 expected gains for $0 \leq p \leq 1$ Correct gains plotted accurately
Solving $8p - 1 = 1 - 4p$	M1		Choosing highest point of their region or correct	
$\Rightarrow p = \frac{1}{6}$	A1			
\Rightarrow Russ plays R_1 with probability $\frac{1}{6}$ and R_2 with prob $\frac{5}{6}$	E1	7		
(ii) Value of game $= \frac{8}{6} - 1$ $= \frac{1}{3}$	B1	1	Or $1 - \frac{4}{6}$	
Total			12	






MD02 (cont)

Q	Solution	Marks	Total	Comments								
4(a)(i)	Slack (variables)	E1	1	Must be correct word								
(ii)	$2x + 2y + z + s = 14$	B1	1	Exactly this								
(b)(i)	Pivot from y -column = 1	B1		Identified or seen used by keeping 3 rd row fixed								
	$\begin{array}{ccccccc c} 1 & -6 & 0 & 5 & 0 & 4 & 0 & 24 \\ 0 & 4 & 0 & -3 & 1 & -2 & 0 & 2 \\ 0 & -1 & 1 & 2 & 0 & 1 & 0 & 6 \\ 0 & 8 & 0 & -5 & 0 & -4 & 1 & 5 \end{array}$	M1		Row operations, even with wrong pivot								
		A1		1st, 2nd or 4th row correct								
		A1	4	All correct								
(ii)	Still negative value in top row	E1	1	(only award if this is true for their tableau)								
(c)(i)	Choosing 4 as pivot in x -column	M1		And perhaps dividing by 4 (using their pivot)								
	$\begin{array}{ccccccc c} 1 & 0 & 0 & \frac{1}{2} & \frac{3}{2} & 1 & 0 & 27 \\ 0 & 1 & 0 & \frac{-3}{4} & \frac{1}{4} & \frac{-1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & 1 & \frac{5}{4} & \frac{1}{4} & \frac{1}{2} & 0 & \frac{13}{2} \\ 0 & 0 & 0 & 1 & -2 & 0 & 1 & 1 \end{array}$	A1		1st, 3rd or 4th row correct ft one slip								
		A1		1st, 3rd or 4th row (another correct) ft one slip								
		A1	4	All correct (condone multiples of rows)								
(ii)	Optimum now reached (since no negatives in top row)	E1		Or maximum value of P indicated (must have no negatives in top row)								
	$P = 27$	B1✓		ft their tableau P								
	$x = \frac{1}{2}, y = 6\frac{1}{2}, z = 0$	B1	3	CAO; final tableau “correct” one slip								
Total			14									
5	July values	B1		3 correct unsimplified								
		B1		Another 3 correct								
		B1		All correct								
	Use of one July min in June calculation	M1										
		A1		4 correct values in June								
		A1		All June values correct (ft one slip)								
	Use of two June min values in May calculation	M1										
		A1		All May correct (ft one slip)								
	Their least May value ⇒ Project for May	M1		Equivalent scheme for Network Method working backwards from August								
	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">May</td> <td style="width: 25%;">June</td> <td style="width: 25%;">July</td> <td style="width: 25%;">August</td> </tr> <tr> <td style="text-align: center;">C</td> <td style="text-align: center;">A</td> <td style="text-align: center;">hol</td> <td style="text-align: center;">B</td> </tr> </table>	May	June	July	August	C	A	hol	B	A1	10	Schedule correct
May	June	July	August									
C	A	hol	B									
				SC B1 if schedule correct with no dynamic programming								

MD02 (cont)

Q	Solution	Marks	Total	Comments																																																																																																																																										
5 (cont)	<table border="1"> <thead> <tr> <th>Stage (Month)</th> <th>State (Projects already done)</th> <th>Action (Project to do)</th> <th>Calculation</th> <th>Cost in thousands of pounds</th> </tr> </thead> <tbody> <tr> <td rowspan="4">August</td> <td><i>A, B, C</i></td> <td>0</td> <td></td> <td>0 (given)</td> </tr> <tr> <td><i>A, B</i></td> <td><i>C</i></td> <td></td> <td>14 (given)</td> </tr> <tr> <td><i>A, C</i></td> <td><i>B</i></td> <td></td> <td>10 (given)</td> </tr> <tr> <td><i>B, C</i></td> <td><i>A</i></td> <td></td> <td>16 (given)</td> </tr> <tr> <td rowspan="12">July</td> <td><i>A, B</i></td> <td>0</td> <td>0 + 14</td> <td>14 (given) ←</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>15 + 0</td> <td>15 (given)</td> </tr> <tr> <td><i>A, C</i></td> <td>0</td> <td>0 + 10</td> <td>10 ←</td> </tr> <tr> <td></td> <td><i>B</i></td> <td>12 + 0</td> <td>12</td> </tr> <tr> <td><i>B, C</i></td> <td>0</td> <td>0 + 16</td> <td>16 ←</td> </tr> <tr> <td></td> <td><i>A</i></td> <td>18 + 0</td> <td>18</td> </tr> <tr> <td><i>A</i></td> <td><i>B</i></td> <td>12 + 14</td> <td>26 (given)</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>15 + 10</td> <td>25 ←</td> </tr> <tr> <td><i>B</i></td> <td><i>A</i></td> <td>18 + 14</td> <td>32</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>15 + 16</td> <td>31 ←</td> </tr> <tr> <td><i>C</i></td> <td><i>A</i></td> <td>18 + 10</td> <td>28 ←</td> </tr> <tr> <td></td> <td><i>B</i></td> <td>12 + 16</td> <td>28 ←</td> </tr> <tr> <td rowspan="12">June</td> <td><i>A</i></td> <td>0</td> <td>0 + 25</td> <td>25 ←</td> </tr> <tr> <td></td> <td><i>B</i></td> <td>13 + 14</td> <td>27</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>17 + 10</td> <td>27</td> </tr> <tr> <td><i>B</i></td> <td>0</td> <td>0 + 31</td> <td>31</td> </tr> <tr> <td></td> <td><i>A</i></td> <td>16 + 14</td> <td>30 ←</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>17 + 16</td> <td>33</td> </tr> <tr> <td><i>C</i></td> <td>0</td> <td>0 + 28</td> <td>28</td> </tr> <tr> <td></td> <td><i>A</i></td> <td>16 + 10</td> <td>26 ←</td> </tr> <tr> <td></td> <td><i>B</i></td> <td>13 + 16</td> <td>29</td> </tr> <tr> <td>0</td> <td><i>A</i></td> <td>16 + 25</td> <td>41 ←</td> </tr> <tr> <td></td> <td><i>B</i></td> <td>13 + 31</td> <td>44</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>17 + 28</td> <td>45</td> </tr> <tr> <td rowspan="4">May</td> <td>0</td> <td>0</td> <td>0 + 41</td> <td>41</td> </tr> <tr> <td></td> <td><i>A</i></td> <td>17 + 25</td> <td>42</td> </tr> <tr> <td></td> <td><i>B</i></td> <td>14 + 30</td> <td>44</td> </tr> <tr> <td></td> <td><i>C</i></td> <td>14 + 26</td> <td>40 ←</td> </tr> </tbody> </table>	Stage (Month)	State (Projects already done)	Action (Project to do)	Calculation	Cost in thousands of pounds	August	<i>A, B, C</i>	0		0 (given)	<i>A, B</i>	<i>C</i>		14 (given)	<i>A, C</i>	<i>B</i>		10 (given)	<i>B, C</i>	<i>A</i>		16 (given)	July	<i>A, B</i>	0	0 + 14	14 (given) ←		<i>C</i>	15 + 0	15 (given)	<i>A, C</i>	0	0 + 10	10 ←		<i>B</i>	12 + 0	12	<i>B, C</i>	0	0 + 16	16 ←		<i>A</i>	18 + 0	18	<i>A</i>	<i>B</i>	12 + 14	26 (given)		<i>C</i>	15 + 10	25 ←	<i>B</i>	<i>A</i>	18 + 14	32		<i>C</i>	15 + 16	31 ←	<i>C</i>	<i>A</i>	18 + 10	28 ←		<i>B</i>	12 + 16	28 ←	June	<i>A</i>	0	0 + 25	25 ←		<i>B</i>	13 + 14	27		<i>C</i>	17 + 10	27	<i>B</i>	0	0 + 31	31		<i>A</i>	16 + 14	30 ←		<i>C</i>	17 + 16	33	<i>C</i>	0	0 + 28	28		<i>A</i>	16 + 10	26 ←		<i>B</i>	13 + 16	29	0	<i>A</i>	16 + 25	41 ←		<i>B</i>	13 + 31	44		<i>C</i>	17 + 28	45	May	0	0	0 + 41	41		<i>A</i>	17 + 25	42		<i>B</i>	14 + 30	44		<i>C</i>	14 + 26	40 ←				
	Stage (Month)	State (Projects already done)	Action (Project to do)	Calculation	Cost in thousands of pounds																																																																																																																																									
	August	<i>A, B, C</i>	0		0 (given)																																																																																																																																									
		<i>A, B</i>	<i>C</i>		14 (given)																																																																																																																																									
		<i>A, C</i>	<i>B</i>		10 (given)																																																																																																																																									
		<i>B, C</i>	<i>A</i>		16 (given)																																																																																																																																									
	July	<i>A, B</i>	0	0 + 14	14 (given) ←																																																																																																																																									
			<i>C</i>	15 + 0	15 (given)																																																																																																																																									
		<i>A, C</i>	0	0 + 10	10 ←																																																																																																																																									
			<i>B</i>	12 + 0	12																																																																																																																																									
		<i>B, C</i>	0	0 + 16	16 ←																																																																																																																																									
			<i>A</i>	18 + 0	18																																																																																																																																									
		<i>A</i>	<i>B</i>	12 + 14	26 (given)																																																																																																																																									
			<i>C</i>	15 + 10	25 ←																																																																																																																																									
		<i>B</i>	<i>A</i>	18 + 14	32																																																																																																																																									
			<i>C</i>	15 + 16	31 ←																																																																																																																																									
		<i>C</i>	<i>A</i>	18 + 10	28 ←																																																																																																																																									
			<i>B</i>	12 + 16	28 ←																																																																																																																																									
	June	<i>A</i>	0	0 + 25	25 ←																																																																																																																																									
			<i>B</i>	13 + 14	27																																																																																																																																									
			<i>C</i>	17 + 10	27																																																																																																																																									
		<i>B</i>	0	0 + 31	31																																																																																																																																									
			<i>A</i>	16 + 14	30 ←																																																																																																																																									
			<i>C</i>	17 + 16	33																																																																																																																																									
		<i>C</i>	0	0 + 28	28																																																																																																																																									
			<i>A</i>	16 + 10	26 ←																																																																																																																																									
			<i>B</i>	13 + 16	29																																																																																																																																									
		0	<i>A</i>	16 + 25	41 ←																																																																																																																																									
			<i>B</i>	13 + 31	44																																																																																																																																									
			<i>C</i>	17 + 28	45																																																																																																																																									
	May	0	0	0 + 41	41																																																																																																																																									
			<i>A</i>	17 + 25	42																																																																																																																																									
			<i>B</i>	14 + 30	44																																																																																																																																									
			<i>C</i>	14 + 26	40 ←																																																																																																																																									
		<table border="1"> <thead> <tr> <th>Schedule</th> <th>May</th> <th>June</th> <th>July</th> <th>August</th> </tr> </thead> <tbody> <tr> <td>Project</td> <td><i>C</i></td> <td><i>A</i></td> <td>holiday</td> <td><i>B</i></td> </tr> </tbody> </table>	Schedule	May	June	July	August	Project	<i>C</i>	<i>A</i>	holiday	<i>B</i>																																																																																																																																		
	Schedule	May	June	July	August																																																																																																																																									
	Project	<i>C</i>	<i>A</i>	holiday	<i>B</i>																																																																																																																																									
		Total		10																																																																																																																																										

MD02 (cont)

Q	Solution	Marks	Total	Comments						
6(a)(i)	Value of cut = $38 + 25 + 0 + 0 + 34$ = 97	B1	1	Must show correct addition AG						
(ii)	$\{S, A\}, \{B, C, T\}$ 65	B1								
	$\{S, B\}, \{A, C, T\}$ 57	B1								
	$\{S, B, C\}, \{A, T\}$ 72	B1								
	$\{S, A, B, C\}, \{T\}$ 56	B1	4							
(iii)	Maximum flow = 53 Minimum cut = Max flow	B1 [✓] E1	2	ft their least cut value						
(iv)	Their max flow on SA, SB or AT, CT All correct AT 22; AC 12; BC 19; CT 31 AB = x; AS = x + 34; SB = 19 - x $0 \leq x \leq 4$	M1 A1	2							
(b)(i)	Initial flow on Figure 6 Forward potential and backward flow Condone 2 slips, ft their Figure 5	M1								
	One correct augmented path in table and correct flow	M1 A1		<table border="1" data-bbox="970 1120 1404 1227"> <thead> <tr> <th>Path</th> <th>Additional Flow</th> </tr> </thead> <tbody> <tr> <td>SBDT</td> <td>6</td> </tr> <tr> <td>SABDCT</td> <td>3</td> </tr> </tbody> </table>	Path	Additional Flow	SBDT	6	SABDCT	3
Path	Additional Flow									
SBDT	6									
SABDCT	3									
	Table correct with total additional flow= 9									
	Final network correct with evidence of labelling procedure used	A1	4							
(ii)	New maximum flow = 62	B1								
	Correct maximum flow on network	B1	2							
	May have 									
	Total		15							
	TOTAL		75							