



---

# A-LEVEL Mathematics

MFP2 Further Pure 2  
Mark scheme

---

6360

June 2017

---

Version: 1.0 Final

---

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [aqa.org.uk](http://aqa.org.uk)

### Key to mark scheme abbreviations

<b>M</b>	mark is for method
<b>dM</b>	mark is dependent on one or more M marks and is for method
<b>A</b>	mark is dependent on M or dM marks and is for accuracy
<b>B</b>	mark is independent of M or dM marks and is for method and accuracy
<b>E</b>	mark is for explanation
<b>FT or ft or F</b>	follow through from previous incorrect result
<b>cao</b>	correct answer only
<b>cso</b>	correct solution only
<b>AWFW</b>	anything which falls within
<b>AWRT</b>	anything which rounds to
<b>ACF</b>	any correct form
<b>AG</b>	answer given
<b>SC</b>	special case
<b>OE</b>	or equivalent
<b>A2,1</b>	2 or 1 (or 0) accuracy marks
<b>-x EE</b>	deduct x marks for each error
<b>NMS</b>	no method shown
<b>PI</b>	possibly implied
<b>SCA</b>	substantially correct approach
<b>c</b>	candidate
<b>sf</b>	significant figure(s)
<b>dp</b>	decimal place(s)

### No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, the principal examiner may suggest that we award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.


**Otherwise we require evidence of a correct method for any marks to be awarded.**

Q 1	Solution	Mark	Total	Comment
<p><b>(a)</b></p> $\frac{A}{2r+1} + \frac{B}{2r+3}$ $A = \frac{1}{4} \quad B = \frac{1}{4}$ <p><b>(b)</b></p> $\frac{A}{3} + \frac{B}{5} - \frac{A}{5} - \frac{B}{7} + \dots$ $[k] \left\{ \frac{1}{3} + (-1)^{n+1} \frac{1}{2n+3} \right\} \text{ OE}$ $\frac{1}{12} + (-1)^{n+1} \frac{1}{4(2n+3)} \text{ OE}$		M1		and attempt to find A or B
		A1	2	$\frac{\frac{1}{4}}{2r+1} + \frac{\frac{1}{4}}{2r+3}$ OE
		M1		clear attempt to use <b>method of differences</b> with “their” A and B
		dM1		condone +, -, ± or $(-1)^n$ instead of $(-1)^{n+1}$ ; may have r for n
		A1	3	must have n
	<b>Total</b>		<b>5</b>	
<p><b>(b)</b></p>	<p>For <b>dM1 correct two</b> remaining terms may be on separate lines with other terms crossed out</p> <p><b>Example 1</b> <math>\frac{1}{3} - \frac{1}{(2n+3)}</math> earns <b>M1 dM1</b></p> <p><b>Example 2</b> <math>\frac{1}{12} \pm \frac{1}{(2n+3)}</math> earns <b>M1 dM0</b></p> <p>Alternative for final <b>A1</b>: n even <math>\frac{1}{12} - \frac{1}{4(2n+3)}</math>; n odd <math>\frac{1}{12} + \frac{1}{4(2n+3)}</math></p>			

Q 2	Solution	Mark	Total	Comment
(a)	$\alpha + \beta + \gamma = -6 + 3i$ <b>OE</b> $\alpha = -3$	<b>M1</b> <b>A1</b>	<b>2</b>	<b>PI</b> by correct $\alpha$
(b) (i)	$\sum \frac{1}{\alpha\beta} = \frac{\alpha + \beta + \gamma}{\alpha\beta\gamma}$ or $i = \frac{\alpha + \beta + \gamma}{\alpha\beta\gamma}$ $q = \frac{6-3i}{i}$ <b>or</b> $\alpha\beta\gamma = 3+6i$ <b>OE</b> $q = -3-6i$ <b>OE</b>	<b>M1</b> <b>A1</b> <b>A1</b>	<b>3</b>	or $\alpha + \beta + \gamma = i \alpha\beta\gamma$ $q$ , $-q$ or $\alpha\beta\gamma$ correct unsimplified <b>PI</b> by correct $q$
(ii)	$-27 + 9(6-3i) - 3p - 3 - 6i = 0$ $p = 8 - 11i$ <b>OE</b>	<b>M1</b> <b>A1</b>	<b>2</b>	correctly substituting “their” values for $\alpha$ and $q$ into equation
(c)	$\sum \alpha^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$ $= 36 - 36i - 9 - 16 + 22i$ $= 11 - 14i$ <b>OE</b>	<b>M1</b> <b>A1cso</b>	<b>2</b>	correct identity
<b>Total</b>			<b>9</b>	
(b)(i)	Withhold final <b>A1</b> if $\alpha + \beta + \gamma = 6 - 3i$ and $\alpha\beta\gamma = q$ leads to correct answer and write <b>FIW</b> Do not treat “1” for “i” as a misread, simply an error			
(b)(ii)	<b>Alternative</b> $\beta\gamma = \frac{\text{“their” } \alpha\beta\gamma}{\alpha} = \text{“their”} - 1 - 2i$ ; $p = \alpha\beta + \beta\gamma + \gamma\alpha = \text{“their” } \alpha(\beta + \gamma) + \beta\gamma$ <b>M1</b> ; $p = 8 - 11i$ <b>A1</b>			
(c)	Withhold <b>A1cso</b> if $\alpha + \beta + \gamma = 6 - 3i$ is seen even if correct answer is given and write <b>FIW</b>			



Q 4	Solution	Mark	Total	Comment
	<p>(a) <math>9k^2 + 17k + 6</math></p> <p>(b) When <math>n = 1</math> <math>LHS = 2</math>; <math>RHS = 2</math> Therefore (formula is) true when <math>n = 1</math></p> <p>Assume result is true for <math>n = k</math> (*) Add <math>(k + 1)</math>th term to both sides</p> $\sum_{r=1}^{k+1} r(2r-1)(3r-1) = \frac{1}{6}k(k+1)(9k^2 - k - 2) + (k+1)(2k+1)(3k+2)$ $= \frac{1}{6}(k+1)\{9k^3 - k^2 - 2k + 6(2k+1)(3k+2)\}$ $\frac{1}{6}(k+1)\{9k^3 + 35k^2 + 40k + 12\}$ $\frac{1}{6}(k+1)(k+2)(9k^2 + 17k + 6)$ $\frac{1}{6}(k+1)(k+2)\{9(k+1)^2 - (k+1) - 2\}$ <p>Hence formula is true for <math>n = k + 1</math> (**) and since true for <math>n = 1</math>, formula is true for <math>n = 1, 2, 3, \dots</math> [by induction] (***)</p>	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>E1</b></p>	<p><b>1</b></p> <p><b>6</b></p> <p><b>7</b></p>	<p>must have this explicit statement</p> <p>adding correct <math>(k+1)</math>th term to RHS</p> <p><b>both</b> sides correct <b>A0</b> if only RHS considered correct quartic is <math>\frac{1}{6}\{9k^4 + 44k^3 + 75k^2 + 52k + 12\}</math> cubic need not have all like terms collected must see this line to earn final <b>A1</b> from part (a)</p> <p>must have (*), (**) and (***) and have earned previous 5 marks</p>
	<b>Total</b>		<b>7</b>	
	<p>(b) For <b>B1</b>, accept “<math>n=1</math> <math>RHS=LHS=2</math>” but must mention here or later that the result is “true when <math>n=1</math>” Do not allow them to simply say “true for all integers <math>n \dots 1</math>” at the end to earn this <b>B1</b> mark. This is <b>B0</b>.</p> <p>Alternative to (***) is “therefore true for all positive integers <math>n</math>” or “so true for all integers <math>n \dots 1</math>” etc However, “true for all <math>n \dots 1</math>” is incorrect and immediately gets <b>E0</b></p> <p>Condone <math>LHS = 1 \times 1 \times 2 + 2 \times 3 \times 5 + \dots + (k+1)(2k+1)(3k+2)</math> <b>OE</b> for first <b>A1</b> but must have “...”</p> <p>May define <math>P(k)</math> as the “proposition that the formula is true when <math>n = k</math>” and earn full marks. However, if <math>P(k)</math> is not defined then allow <b>B1</b> for showing <math>P(1)</math> is true but withhold <b>E1</b> mark.</p>			

Q 5	Solution	Mark	Total	Comment	
<b>(a)(i)</b>	$\tan^{-1} \frac{3}{\sqrt{3}} = \frac{\pi}{3}$	<b>M1</b>		or finding angle to Im(z) axis = $\frac{\pi}{6}$	
	$(\arg \omega =) \frac{2\pi}{3}$	<b>A1</b>	<b>2</b>	<b>PI</b> by correct answer	
	<b>(ii)</b>	$( \omega - 2i ^2) = (-\sqrt{3})^2 + 1^2$	<b>M1</b>		<b>PI</b> by correct answer
		$ \omega - 2i  = 2$	<b>A1</b>	<b>2</b>	
	<b>(b)(i)</b>		<b>M1</b>		arc of circle in second quadrant
			<b>A1</b>		circle centre at 2i (2 marked on Im(z)-axis) and touching real axis at O
			<b>M1</b>		line from O to at least edge of circle
			<b>A1</b>		inclined at roughly $\frac{\pi}{3}$ to negative real axis as drawn
			<b>A1</b>	<b>5</b>	must have earned previous 4 marks correct shading of region bounded by line, imaginary axis and circular arc
	<b>(ii)</b>	$\omega$ marked correctly	<b>B1</b>	<b>1</b>	clear intention to be at intersection point <b>B0</b> if only line or circle drawn
<b>(iii)</b>	Max value = $\left  \frac{\omega}{2} - 4i \right $ or $\left  -\frac{\sqrt{3}}{2} + \frac{3i}{2} - 4i \right $	<b>M1</b>		correct expression- distance from $\frac{1}{2}\omega$ to 4i that could be evaluated to give correct ans	
	or $\sqrt{\frac{ \omega ^2}{4} + 2^2}$ etc <b>ACF</b> $= \sqrt{7}$	<b>A1</b>	<b>2</b>	$d^2 = \frac{ \omega ^2}{4} + 4^2 - 4 \omega  \cos \frac{\pi}{6}$	
<b>Total</b>			<b>12</b>		
<b>(a)(i)</b>	NMS $(\theta =) \frac{2\pi}{3}$ <b>M1 A1</b> ; $\tan^{-1} \left( -\frac{3}{\sqrt{3}} \right) = -\frac{\pi}{3}$ or sight of $-\frac{\pi}{3}$ earns <b>M1</b>				
<b>(b)(i)</b>	Allow freehand circle and clear “intention” to touch real axis at origin First <b>A1</b> : condone 2i or (0,2) or 2 clear dashes to indicate centre on Im(z) axis or radius indicated as 2 <b>and</b> circle touching real axis at O Second <b>A1</b> : award if angle made with negative Re(z) axis is greater than $\frac{\pi}{4}$				
<b>(iii)</b>	Condone circle/line as dotted lines NMS max = $\sqrt{7}$ scores full marks				



Q 6	Solution	Mark	Total	Comment
	$\frac{x^2}{2} \tan^{-1}\left(\frac{x}{\sqrt{3}}\right) - \int k x^2 \frac{1}{1 + \left(\frac{x}{\sqrt{3}}\right)^2} (dx)$ $\frac{x^2}{2} \tan^{-1}\left(\frac{x}{\sqrt{3}}\right) - \int \frac{x^2}{2} \times \frac{1}{\sqrt{3}} \times \frac{1}{1 + \frac{x^2}{3}} (dx)$ $\frac{x^2}{x^2 + A} = 1 - \frac{A}{x^2 + A} \quad \text{OE}$ $\frac{x^2}{2} \tan^{-1}\left(\frac{x}{\sqrt{3}}\right) - \frac{\sqrt{3}}{2} x + \frac{3}{2\sqrt{3}} \times \sqrt{3} \times \tan^{-1}\left(\frac{x}{\sqrt{3}}\right)$ $I = \frac{(\sqrt{3})^2}{2} \tan^{-1}(1) - \frac{\sqrt{3}}{2} \sqrt{3} + \frac{3}{2\sqrt{3}} \times \sqrt{3} \times \tan^{-1}(1)$ $= \frac{3\pi}{8} - \frac{3}{2} + \frac{3\pi}{8}$ $= \frac{3\pi}{4} - \frac{3}{2}$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>B1F</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p><b>6</b></p>	<p>Integration by parts – at least this far – (denominator may be <math>3 + x^2</math>)</p> <p>or <math>\frac{x^2}{2} \tan^{-1}\left(\frac{x}{\sqrt{3}}\right) - \int \frac{x^2}{2} \times \frac{\sqrt{3}}{3 + x^2} (dx)</math> <b>OE</b></p> <p>or <math>\frac{x^2}{1 + \frac{x^2}{3}} = 3 - \frac{3}{1 + \frac{x^2}{3}}</math> etc</p> <p>correct unsimplified</p> <p>correct unsimplified sub of limits</p>
	<b>Total</b>		<b>6</b>	
	<p>Do NOT allow misread of <math>\frac{x}{3}</math> for <math>\frac{x}{\sqrt{3}}</math>; it eases the question considerably</p> <p><b>Alternative 1:</b> <math>x = \sqrt{3} u</math>; <math>I = \int 3u \tan^{-1} u \, du = \frac{3}{2} u^2 \tan^{-1} u - k \int \frac{u^2}{1 + u^2} \, du</math> <b>M1</b>; <math>k = \frac{3}{2}</math> <b>A1</b>; <math>\frac{u^2}{1 + u^2} = 1 - \frac{1}{1 + u^2}</math> <b>B1</b>;</p> <p><math>\frac{3u^2}{2} \tan^{-1} u - \frac{3}{2} u + \frac{3}{2} \tan^{-1} u</math> <b>A1</b>; then <b>A1 A1</b> as above.</p> <p><b>Alternative 2:</b> <math>x = \sqrt{3} \tan u</math>; <math>\frac{dx}{du} = \sqrt{3} \sec^2 u</math>; <math>I = \int 3u \tan u \sec^2 u \, du</math></p> <p><math>I = \frac{3}{2} [u \tan^2 u] - \int k \tan^2 u \, du</math> <b>M1</b> <math>k = \frac{3}{2}</math> <b>A1</b> (correct)</p> <p>replacing <math>\tan^2 u</math> by <math>\sec^2 u - 1</math> in integral <b>dM1</b>; <math>I = \frac{3}{2} [u \tan^2 u] + \frac{3}{2} u - \frac{3}{2} \tan u</math> <b>A1</b>; then <b>A1 A1</b> as above.</p>			

Q 7	Solution	Mark	Total	Comment
(a)	$\frac{(1 + \cosh \theta) \cosh \theta - \sinh \theta \sinh \theta}{(1 + \cosh \theta)^2}$ Numerator = $\cosh \theta + 1$ $\times \frac{1 + \cosh \theta}{\sinh \theta}$ $f'(\theta) = \frac{1}{\sinh \theta}$	<b>M1</b>  <b>A1</b>  <b>dM1</b>  <b>A1</b>	<b>4</b>	quotient rule correct  correctly simplified  <b>AG</b> – no errors seen and $f'(\theta) = \dots$
(b)(i)	$\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right] = 1 + \left( \frac{1}{x} \right)^2$ $\frac{x^2 + 1}{x^2} \quad \text{or} \quad \sqrt{\frac{x^2 + 1}{x^2}} \quad \text{or} \quad \frac{\sqrt{x^2 + 1}}{\sqrt{x^2}}$ $s = \int_1^{2\sqrt{2}} \frac{\sqrt{x^2 + 1}}{x} dx$	<b>M1</b>  <b>A1</b>  <b>A1</b>	<b>3</b>	condone $\int \sqrt{1 + \left( \frac{1}{x} \right)^2} (dx)$ for <b>M1</b>  Allow this mark but withhold final <b>A1</b> mark if $\frac{dy}{dx}$ or $\left( \frac{dy}{dx} \right)^2$ not seen  <b>AG</b> (be convinced) - must have “s =”, limits and dx and must have $\sqrt{x^2 + 1}$ in numerator
(ii)	$x = \sinh \theta \quad dx = \cosh \theta d\theta$ <b>OE</b> $\int \frac{\cosh \theta \cosh \theta}{\sinh \theta} d\theta \quad (\text{must have } d\theta)$ Attempt to split into two terms using $\cosh^2 \theta = \pm 1 \pm \sinh^2 \theta$ $\int \left( \frac{1}{\sinh \theta} + \sinh \theta \right) [d\theta]$ $\ln \left[ \frac{\sinh \theta}{1 + \cosh \theta} \right] + \cosh \theta$ $s = \ln \left[ \frac{2\sqrt{2}}{1+3} \right] + 3 - \ln \left[ \frac{1}{1+\sqrt{2}} \right] - \sqrt{2}$ $3 - \sqrt{2} + \ln \left( 1 + \frac{\sqrt{2}}{2} \right)$	<b>M1</b>  <b>A1</b>  <b>M1</b>  <b>A1</b>  <b>dM1</b>  <b>A1</b>  <b>A1</b>	<b>7</b>	or $x = \tan \theta \quad dx = \sec^2 \theta d\theta$ $\int \frac{\sec \theta \sec^2 \theta}{\tan \theta} d\theta$  <b>PI</b> by correct split below  correct split and must have integral sign  integrating $\pm \frac{1}{\sinh \theta} \pm \sinh \theta$ correctly  correct unsimplified  <b>AG</b> partly so be convinced
<b>Total</b>			<b>13</b>	
(a)	<b>Alternative:</b> $[f(\theta) =] \ln(\sinh \theta) - \ln(1 + \cosh \theta)$ <b>and</b> one term differentiated correctly <b>M1</b> $[f'(\theta) =] \frac{\cosh \theta}{\sinh \theta} - \frac{\sinh \theta}{1 + \cosh \theta} \quad \mathbf{A1} = \frac{(1 + \cosh \theta) \cosh \theta - \sinh^2 \theta}{\sinh \theta (1 + \cosh \theta)} \quad \mathbf{dM1}$ (common denominator) $f'(\theta) = \frac{1}{\sinh \theta} \quad \mathbf{A1}$ ( <b>AG</b> no errors seen and $f'(\theta) = \dots$ )			
(b)(ii)	In alternative on RHS; <b>B1</b> for using $\sec^2 \theta = 1 + \tan^2 \theta$ used in numerator; <b>dM1</b> for splitting integrand $\pm \frac{1}{\sin \theta} \pm \sec \theta \tan \theta$ and <b>dM1</b> for integrating correctly <b>NB</b> $\int (\csc \theta + \sec \theta \tan \theta) d\theta = -\ln(\csc \theta + \cot \theta) + \sec \theta$			

Q 8	Solution	Mark	Total	Comment	
<b>(a)</b>	$\cos 7\theta + i \sin 7\theta = (\cos \theta + i \sin \theta)^7$	<b>B1</b>		or $\sin 7\theta = \text{Im part of } (\cos \theta + i \sin \theta)^7$ <b>PI</b> by later work	
	$[c^7] + 7c^6(is) + [21c^5(is)^2] + 35c^4(is)^3$	<b>M1</b>		condone up to 2 errors in imaginary part of expansion for <b>M1</b> – ignore real terms	
	$[+35c^3(is)^4] + 21c^2(is)^5 + [7c(is)^6] + (is)^7$	<b>A1</b>		correct imaginary terms	
	$7(1-s^2)^3(is) + 35(1-s^2)^2(is)^3$	<b>dM1</b>		correct use of $c^2 = 1-s^2$ in at least two imaginary terms – i.e. $c^6 = (1-s^2)^3$ etc	
	$+ 21(1-s^2)(is)^5 + (is)^7$				
	$\sin 7\theta = 7s(1-3s^2+3s^4-s^6)$	<b>A1</b>		RHS correct unsimplified expansion <b>and</b> equated to $\sin 7\theta$	
	$-35s^3(1-2s^2+s^4) + 21s^5(1-s^2) - s^7$				
	$\frac{\sin 7\theta}{\sin \theta} = 7 - 56\sin^2 \theta + 112\sin^4 \theta - 64\sin^6 \theta$	<b>A1</b>	<b>6</b>	<b>AG</b> be convinced – terms must be in this order	
	<b>(b)(i)</b>	$\sin 7\theta = 0 \Rightarrow 7\theta = (n)\pi \Rightarrow \theta = (n)\frac{\pi}{7}$	<b>M1</b>		condone no mention of “but $\sin \theta \neq 0$ ”
		$x = \sin^2 \theta$ seen or used	<b>E1</b>		must earn <b>M1</b> and have/use $x = \sin^2 \theta$ <b>and</b> statement
so $\sin^2 \frac{\pi}{7}$ is a root of cubic equation					
other roots are $\sin^2 \frac{2\pi}{7}$ & $\sin^2 \frac{3\pi}{7}$ <b>OE</b>	<b>B1</b>	<b>3</b>	accept $\sin^2 \frac{4\pi}{7}$ & $\sin^2 \frac{5\pi}{7}$ etc but <b>not</b> $\sin^2 \frac{3\pi}{7}$ & $\sin^2 \frac{4\pi}{7}$ etc		
<b>(ii)</b>	Considering $\sum \frac{1}{\alpha}$	<b>M1</b>		must relate $\sin^2 \frac{\pi}{7}$ etc to $\alpha, \beta, \gamma$	
	$\frac{\alpha\beta + \beta\gamma + \gamma\alpha}{\alpha\beta\gamma}$	<b>A1</b>			
	$= \frac{56/64}{7/64} = 8$	<b>A1</b>	<b>3</b>	do <b>not</b> accept $\frac{56}{7}$ if using this approach	
<b>Total</b>			<b>12</b>		
<b>(b)(i)</b>	Condone reverse argument namely $\theta = \frac{\pi}{7} \Rightarrow \sin 7\theta = 0$ for <b>M1</b>				
	$\frac{\pi}{7}$ is a root of $\frac{\sin 7\theta}{\sin \theta} = 0$ earns <b>M1</b>				
<b>(ii)</b>	<b>Alternative</b> : put $z = 1/y$ <b>M1</b>				
	new equation $7z^3 - 56z^2 + 112z - 64 = 0$ <b>A1</b> ; sum of these roots $= \frac{56}{7} = 8$ <b>A1</b>				
	<b>NMS</b> 8 scores no marks				