



**General Certificate of Education**

**Mathematics 6360**

**MPC4      Pure Core 4**

**Mark Scheme**

*2010 examination - January series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: [www.aqa.org.uk](http://www.aqa.org.uk)

Copyright © 2010 AQA and its licensors. All rights reserved.

#### COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

---

**Key to mark scheme and abbreviations used in marking**

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

**No Method Shown**

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

Q	Solution	Marks	Total	Comments
<b>1(a)(i)</b> <b>(ii)</b>  <b>(b)</b>	$f(-1) = -15 + 19 - 4 = 0$	B1	1	evaluate <b>or</b> complete division leading to a numerical remainder  Or decimal equivalent $(0.96 + 3.04 - 4)$ <b>or</b> zero remainder $\Rightarrow$ factor  Stated or implied.  Any appropriate method to find third factor  $\left. \begin{array}{l} (5x-2)(3x^2 \pm 5x \pm 2) + \text{attempt} \\ \text{to factorise} \\ \text{Factorise numerator correctly} \\ \text{and attempt to simplify} \end{array} \right\}$  CSO no ISW
	$f\left(\frac{2}{5}\right)$	M1		
	$\left(15 \times \frac{8}{125} + 19 \times \frac{4}{25} - 4\right) = 0 \Rightarrow$ factor	A1	2	
	$(x+1)$ is a factor	B1		
	Third factor is $(3x+2)$	M1 A1		
	$\frac{15x^2 - 6x}{f(x)} = \frac{3x(5x-2)}{(x+1)(5x-2)(3x+2)}$	M1		
	$= \frac{3x}{(x+1)(3x+2)}$	A1	5	
	<b>Total</b>		<b>8</b>	
<b>2(a)</b>  <b>(b)(i)</b>  <b>(ii)</b>  <b>(c)</b>	$R = \sqrt{10}$ $\tan \alpha = 3$ $\alpha = 1.249$ ignore extra out of range	B1 M1 A1	3	Accept $R = 3.16$ or better OE AWRT 1.25 SC $\alpha = 0.322$ B1 radians only  F on $R$  AWRT 4.39 $51.56^\circ$ or .. $57^\circ$ or better  Two values, accept 2dp and condone 5.4 condone use of degrees  F on $x - \alpha$ , either value. AWRT  CSO 3dp or better
	minimum value $= -\sqrt{10}$	B1F	1	
	$\cos(x - \alpha) = -1$ $x = 4.391$	M1 A1F	2	
	$\cos(x - \alpha) = \frac{2}{\sqrt{10}}$ $x - \alpha = \pm 0.886$ 5.397 ignore extra out of range	M1 A1		
	$x = 0.36296..$ 2.13512.. $x = 0.363$ 2.135	A1F A1	4	
	<b>Total</b>		<b>10</b>	
<b>(c)</b>	<b>Alternative</b> $10 \sin^2 x - 12 \sin x + 3 = 0$  $\sin x =$ two numerical answers $-1 \leq \text{ans} \leq 1$  $x =$ one correct answer  $x = 0.363$ 2.135	M1 A1F A1F A1		Or equivalent quadratic using $\cos x$ (ie $\sin^2 x + \cos^2 x = 1$ used) Or equivalent using $\cos x$  CSO 3 dp or better

**MPC4 (cont)**

<b>Q</b>	<b>Solution</b>	<b>Marks</b>	<b>Total</b>	<b>Comments</b>
<b>3(a)(i)</b>	$(1+x)^{\frac{1}{3}} = 1 \pm \frac{1}{3}x + kx^2$ $= 1 - \frac{1}{3}x + \frac{2}{9}x^2$	M1 A1	2	$1 \pm \frac{1}{3}x + kx^2$
<b>(ii)</b>	$\left(1 + \frac{3}{4}x\right)^{\frac{1}{3}} = 1 - \frac{1}{3} \times \frac{3}{4}x + \frac{2}{9}\left(\frac{3}{4}x\right)^2$ $= 1 - \frac{1}{4}x + \frac{1}{8}x^2$	M1 A1	2	$x$ replaced by $\frac{3}{4}x$ or start binomial again; condone missing brackets
<b>(b)</b>	$\sqrt[3]{\frac{256}{4+3x}} = k \left(1 + \frac{3}{4}x\right)^{\frac{1}{3}}$ $= 4 \left(1 - \frac{1}{4}x + \frac{1}{8}x^2\right)$ $= 4 - x + \frac{1}{2}x^2 \quad \text{or}$ $a = 4 \quad b = -1 \quad c = \frac{1}{2}$	M1 A1F A1	3	$k \neq 1$ F on (a)(ii) $k = 4$ , accept $\sqrt[3]{64}$ or $64^{\frac{1}{3}}$ CSO fully simplified Be convinced
	<b>Total</b>		<b>7</b>	
<b>4(a)</b>	$10x^2 + 8 = 2(x+1)(5x-1) +$ $A(5x-1) + B(x+1)$ $x = -1 \quad x = \frac{1}{5}$ $A = -3 \quad B = 7$	M1 A1 m1 A1	4	$A$ and $B$ terms correct  Use two values of $x$ to find $A$ and $B$ , or set up and solve $8 + 5A + B = 0$ $-2 - A + B = 8$ SC1 NWS $A$ & $B$ correct $\frac{4}{4}$ SC2 NWS $A$ or $B$ correct $\frac{1}{4}$
<b>(b)</b>	$\int \frac{10x^2 + 8}{(x+1)(5x-1)} dx = \int 2 - \frac{3}{x+1} + \frac{7}{5x-1} dx$ $= 2x + C$ $-3 \ln(x+1) + \frac{7}{5} \ln(5x-1)$	M1  B1 M1  A1F	4	Use the partial fractions  $a \ln(x+1) + b \ln(5x-1)$ condone missing brackets F on $A$ and $B$
	<b>Total</b>		<b>8</b>	
<b>5</b>	$x^2 + xy = e^y$ $2x + y + x \frac{dy}{dx} = e^y \frac{dy}{dx}$ $(-1, 0) \quad \frac{dy}{dx} = -1$	B1 M1 A1 B1  A1	5	$2x$ Use product rule  RHS  CSO
	<b>Total</b>		<b>5</b>	

## MPC4 (cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	$\sin 2\theta = 2 \sin \theta \cos \theta$ $\cos 2\theta = \cos^2 \theta - \sin^2 \theta$	B1 B1	2	OE condone use of $x$ etc, but variable must be consistent
(ii)	$\sin \theta = \frac{4}{5} \Rightarrow \sin 2\theta = 2 \times \frac{4}{5} \times \frac{3}{5} = \frac{24}{25}$ <b>or</b> $2 \times \sin \left( \cos^{-1} \frac{3}{5} \right) \times \frac{3}{5}$ $\cos 2\theta = \frac{9}{25} - \frac{16}{25} = -\frac{7}{25}$	B1		AG Use of $106.26^\circ \dots$ B0
(b)(i)	$\frac{dx}{d\theta} = 6 \cos 2\theta$ , $\frac{dy}{d\theta} = -8 \sin 2\theta$ $\frac{dy}{dx} = -\frac{4 \sin 2\theta}{3 \cos 2\theta}$ ISW	M1 A1 A1	3	Attempt both derivatives. ie $p \cos 2\theta$ Both correct. $q \sin 2\theta$ CSO OE
(ii)	$P \left( \frac{72}{25}, -\frac{28}{25} \right)$ Gradient = $-\frac{4}{3} \times -\frac{24}{7}$	B1F M1		(2.88, -1.12) Their $\frac{q \sin 2\theta}{p \cos 2\theta}$ <b>or</b> $\frac{p \cos 2\theta}{q \sin 2\theta}$ must be working with rational numbers
	Tangent $y + \frac{28}{25} = \frac{32}{7} \left( x - \frac{72}{25} \right)$ ISW	A1	3	Any correct form. $7y = 32x - 100$ Fractions in simplest form Equation required
<b>Total</b>			<b>10</b>	

## MPC4 (cont)

Q	Solution	Marks	Total	Comments
7	$\int y \, dy = \int \cos\left(\frac{x}{3}\right) dx$ $\frac{1}{2}y^2 = 3\sin\left(\frac{x}{3}\right) + C$ $\left(\frac{\pi}{2}, 1\right) \quad \frac{1}{2} = 3\sin\frac{\pi}{6} + C$ $C = -1$ $y^2 = 6\sin\left(\frac{x}{3}\right) - 2$	<p>B1</p> <p>B1 B1</p> <p>M1</p> <p>A1F</p> <p>A1</p>	6	<p>Separate; condone missing integral signs.</p> <p>Accept <math>\frac{\sin\left(\frac{x}{3}\right)}{\frac{1}{3}}</math></p> <p>Use <math>\left(\frac{\pi}{2}, 1\right)</math> to find C</p> <p>must be in form <math>py^2 = q\sin\left(\frac{x}{3}\right) + C</math></p> <p>CSO</p>
<b>Total</b>			<b>6</b>	
<b>8(a)</b>	$0 = 2 + \lambda \Rightarrow \lambda = -2$ <b>Check</b> $-1 + -2 \times -3 = -1 + 6 = 5$ $-5 - 2 \times 2 = -5 \times -4 = -9$	<p>M1</p> <p>A1</p>	2	OE
<b>(b)</b>	$\overrightarrow{BC} = \begin{bmatrix} 9 \\ 2 \\ 3 \end{bmatrix} - \begin{bmatrix} 0 \\ 5 \\ -9 \end{bmatrix} = \begin{bmatrix} 9 \\ -3 \\ 12 \end{bmatrix}$	<p>M1</p> <p>A1</p>	2	$\pm (\overrightarrow{OC} - \overrightarrow{OB})$
<b>(c)(i)</b>	$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{AD} = \overrightarrow{OA} + 2\overrightarrow{BC}$ $\overrightarrow{OD} = \begin{bmatrix} 2 \\ -1 \\ -5 \end{bmatrix} + \begin{bmatrix} 18 \\ -6 \\ 24 \end{bmatrix} = \begin{bmatrix} 20 \\ -7 \\ 19 \end{bmatrix}$ <p>D is (20, -7, 19)</p>	<p>M1</p> <p>A1</p>	2	AG
<b>(ii)</b>	$\overrightarrow{PD} = \overrightarrow{OD} - \overrightarrow{OP} =$ $\begin{bmatrix} 20 \\ -7 \\ 19 \end{bmatrix} - \begin{bmatrix} 2+p \\ -1-3p \\ -5+2p \end{bmatrix} = \begin{bmatrix} 18-p \\ -6+3p \\ 24-2p \end{bmatrix}$ $\overrightarrow{PD} \cdot \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix} = 0$ $(18-p) \times 1 + (-6+3p) \times -3 + (24-2p) \times 2 = 0$ $p = 6$	<p>M1</p> <p>A1</p> <p>B1</p> <p>m1</p> <p>A1</p>	5	<p>Find <math>\overrightarrow{PD}</math> in terms of <math>p</math></p> <p>condone <math>\overrightarrow{PD} = \overrightarrow{OP} - \overrightarrow{OD}</math> here</p> <p>CSO OE working with <math>\overrightarrow{DP}</math></p>
<b>Total</b>			<b>11</b>	

## MPC4 (cont)

Q	Solution	Marks	Total	Comments
9(a)(i)	$t = 0 \quad h = A(1 - 1) = 0$	B1	1	
(ii)	$57 = A \left( 1 - e^{-\frac{12}{4}} \right)$	M1		
	$A = \frac{57}{(1 - e^{-3})} \approx 60$	A1	2	Or 59.9... seen. $A = \text{correct expression} \approx 60 \text{ 2sf}$
(b)(i)	$h = 48 \quad \frac{48}{60} = 1 - e^{-\frac{1}{4}t}$	M1		
	$\ln \left( e^{-\frac{1}{4}t} \right) = \ln \left( \frac{1}{5} \right)$	m1		
	$-\frac{1}{4}t = -\ln 5 \Rightarrow t = 4 \ln 5$	A1	3	
(ii)	$\frac{dh}{dt} = -\frac{1}{4} \times -60 \times e^{-\frac{1}{4}t}$	M1		Differentiate, condone sign errors
	$60e^{-\frac{1}{4}t} = 60 - h \Rightarrow \frac{dh}{dt} = \frac{1}{4}(60 - h)$	m1		Eliminate $e^{-\frac{1}{4}t}$
	$\frac{dh}{dt} = 15 - \frac{h}{4}$	A1	3	CSO, AG
(iii)	$h = 8$	B1	1	
	<b>Total</b>		<b>10</b>	
	<b>TOTAL</b>		<b>75</b>	